

FORM PTO-1350
(REV 10-94)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEYS DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

270.62USWO

U.S. APPLICATION NO. (If known, use 37 C.F.R. 1.5)

Unknown **097 980364**

INTERNATIONAL APPLICATION NO.

PCT/CA00/00642

INTERNATIONAL FILING DATE

June 2, 2000

PRIORITY DATE CLAIMED

June 2, 1999

TITLE OF INVENTION

USE OF THE BNM3 TRANSCRIPTIONAL ACTIVATOR TO CONTROL PLANT EMBRYOGENESIS AND REGENERATION
PROCESS

APPLICANT(S) FOR DO/EO/US

BOUTILIER et al.

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(I).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US)
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An unsigned oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern document(s) or information included:

11. ☒ An Information Disclosure Statement under 37 CFR 1.97 and 1.98, form 1449, 10 references.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A **FIRST** preliminary amendment.
☐ A **SECOND** of **SUBSEQUENT** preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information: International Search Report; International Preliminary Examination Report; PCT Request; PCT/IPEA/402; PCT/IB/306; PCT/IB/332; PCT/IB/304; PCT/IPEA/408; 13 pages of Sequence Listing; copy of published PCT application

U.S. APPLICATION NO. (if known, see 37 C.F.R. 1.2) Unknown 091/980364	INTERNATIONAL APPLICATION NO PCT/CA00/00642	ATTORNEY'S DOCKET NUMBER 270.62USWO
--	--	--

17. <input checked="" type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492(a) (1)-(5)): Search Report has been prepared by the EPO or JPO.....\$890.00 International preliminary examination fee paid to USPTO (37 CFR 1.492(a)(1)).....\$710.00 No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2)).....\$740.00 Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(3)) paid to USPTO \$1040.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4)\$100.00	CALCULATIONS PTO USE ONLY
---	---

ENTER APPROPRIATE BASIC FEE AMOUNT =		\$890.00	
Surcharge of \$130.00 for furnishing the oath or declaration later than [] 20 [] 30 months from the earliest claimed priority date (37 CFR 1.492(e)).		\$0	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE
Total claims	74 -20 =	54	X \$18.00
Independent claims	4 -3 =	1	X \$84.00
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$260.00
TOTAL OF ABOVE CALCULATIONS =		\$1946.00	
Reduction by 1/2 for filing by small entity, if applicable. Small entity status is claimed pursuant to 37 CFR 1.27		\$0	
SUBTOTAL =		\$1946.00	
Processing fee of \$130.00 for furnishing the English translation later than [] 20 [] 30 months from the earliest claimed priority date (37 CFR 1.492(f)).		+	\$0
TOTAL NATIONAL FEE =		\$1946.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property		+	\$0
TOTAL FEES ENCLOSED =		\$1946.00	
		Amount to be: refunded	\$0
		charged	\$0

a. ☒ Check in the amount of \$1946.00 to cover the above fees is enclosed.

b. ☐ Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees.
 A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any
 overpayment to Deposit Account No. 13-2725.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO Gregory A. Sebald MERCHANT & GOULD P.O. Box 2903 Minneapolis, MN 55402-0903	SIGNATURE: NAME: Gregory A. Sebald REGISTRATION NUMBER: 33,280
--	--

S/N unknown

PATENTIN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant:	BOUTILIER et al.	Docket No.:	270.62USWO
Serial No.:	unknown	Filed:	concurrent herewith
Int'l Appln No.:	PCT/CA00/00642	Int'l Filing Date:	June 2, 2000
Title:	USE OF THE BNM3 TRANSCRIPTIONAL ACTIVATOR TO CONTROL PLANT EMBRYOGENESIS AND REGENERATION PROCESS		

CERTIFICATE UNDER 37 CFR 1.10

'Express Mail' mailing label number: EV 037641387 US

Date of Deposit: November 30, 2001

I hereby certify that this correspondence is being deposited with the United States Postal Service 'Express Mail Post Office To Addressee' service under 37 CFR 1.10 on the date indicated above and is addressed to: Box PCT, Assistant Commissioner for Patents, P.O. Box 2327, Arlington, VA 22202.

By: 

Name: Chris Stordahl

PRELIMINARY AMENDMENT

Box PCT
Assistant Commissioner for Patents
P.O. Box 2327
Arlington, VA 22202

Dear Sir:

In connection with the above-identified application filed herewith, please enter the following preliminary amendment (marked-up copy attached):

IN THE ABSTRACT

Insert the attached Abstract page into the application as the last page thereof.

IN THE SPECIFICATION

A courtesy copy of the present specification is enclosed herewith. However, the World Intellectual Property Office (WIPO) copy should be relied upon if it is already in the U.S. Patent Office.

200011-11500000

IN THE CLAIMS

Please amend the following claims:

10. (Amended) A vector comprising the isolated DNA molecule as claimed in claim 1, wherein said isolated DNA molecule is under control of a regulatory element that directs expression of said DNA in a plant cell.

15. (Amended) A transformed plant cell comprising the vector of claim 10.

16. (Amended) A transformed plant comprising the vector of claim 10.

18. (Amended) An isolated protein encoded by the isolated DNA molecule as claimed in claim 4.

19. (Amended) A method of producing asexually derived embryos comprising:

- i) transforming a plant cell with the vector of claim 10;
- ii) growing said plant cell to produce transformed tissue;
- iii) selecting said transformed tissue for occurrence of said isolated DNA molecule; and
- iv) assaying said transformed plant for asexual embryo production.

25. (Amended) A method of modifying the regenerative capacity of a plant comprising:

- i) transforming a plant cell with the vector of claim 10;
- ii) growing said transformed plant cell to produce transformed tissue; and
- iii) assaying said transformed plant tissue for enhanced regeneration as compared to wild-type tissue.

27. (Amended) A method of selecting a transformed plant comprising;

- i) transforming a normally non-regenerative plant with a vector of claim 10; and

- ii) determining whether said transformed plant is able to regenerate under conditions in which said normally non-regenerative plant does not regenerate.

30. (Amended) A vector comprising the isolated DNA molecule of claim 28 operably associated with a gene of interest, wherein said isolated DNA molecule directs the expression of said gene of interest within a plant cell.

33. (Amended) A transformed plant cell comprising the vector of claim 30.

34. (Amended) A transformed plant comprising the vector of claim 30.

36. (Amended) A method for directing the expression of a gene of interest within a developing embryo of a plant comprising transforming said plant with the vector as defined by claim 30.

37. (Amended) A use of a nucleotide sequence as defined in claim 4 as a selectable marker.

38. (Amended) A method of producing asexually derived embryos comprising:

- i) transiently transforming a plant cell with the vector of claim 10, to produce a modified plant cell;
- ii) growing said modified plant cell to produce tissue; and
- iii) assaying said tissue for asexual embryo formation.

44. (Amended) A method of modifying the regenerative capacity of a plant comprising

- i) transiently transforming a plant cell with the vector of claim 10, to produce a modified plant cell;
- ii) growing said modified plant cell to produce tissue; and
- iii) assaying said tissue for enhanced regeneration as compared to wild-type tissue.

46. (Amended) A method of producing an apomictic plant comprising:

- i) transforming a plant with the vector of claim 10, to produce a transformed plant;
- ii) selecting said transformed plant for occurrence of said isolated DNA molecule; and
- iii) assaying said transformed plant for asexual embryo production.

51. (Amended) A method of modifying the regenerative capacity of a plant comprising

- i) transiently transforming a plant cell with the vector of claim 10;
- ii) growing said plant cell to form tissue; and
- iii) assaying said tissue for enhanced regeneration as compared to wild-type tissue.

53. (Amended) A method of selecting a modified plant comprising;

- i) transiently transforming a normally non-regenerative plant with a vector of claim 10 to produce said modified plant; and
- ii) determining whether said modified plant is able to regenerate under conditions in which said normally non-regenerative plant does not germinate.

55. (Amended) A method of producing a protein of interest comprising

- i) transforming a plant with a vector of claim 10 to produce a transformed plant;
- ii) selecting said transformed plant for occurrence of said isolated DNA molecule; and
- iv) growing said transformed plant in order to produce said protein of interest, wherein expression of said protein of interest is induced by the expression product of said isolated DNA.

58. (Amended) The method of claim 55, wherein said protein of interest is selected from the group consisting of a pharmaceutically active protein, antibody, industrial enzyme, protein supplement, nutraceutical, storage protein, an enzyme involved in oil biosynthesis, animal feed, and animal feed supplement.

59. (Amended) The isolated DNA molecule of claim 4, wherein said isolated DNA molecule encodes a protein that is at least 70% similar with the amino acid defined by SEQ ID NO:2.

60. (Amended) The isolated DNA molecule of claim 4, wherein said isolated DNA molecule encodes a protein that is at least 70% similar with the amino acid defined by SEQ ID NO:4.

66. (Amended) A vector comprising the isolated DNA molecule as claimed in claim 63, wherein said isolated DNA molecule is under control of a regulatory element that directs expression of said DNA in a plant cell.

71. (Amended) A transformed plant cell comprising the vector of claim 66.

72. (Amended) A transformed plant comprising the vector of claim 66.

REMARKS

The above preliminary amendment is made to remove multiple dependencies from claims 10, 15, 16, 18, 19, 25, 27, 30, 33, 34, 36, 37, 38, 44, 46, 51, 53, 55, 58, 59, 60, 66, 71 and 72.

A new abstract page is supplied to conform to that appearing on the publication page of the WIPO application, but the new Abstract is typed on a separate page as required by U.S. practice.


Applicants respectfully request that the preliminary amendment described herein be entered into the record prior to calculation of the filing fee and prior to examination and consideration of the above-identified application.

If a telephone conference would be helpful in resolving any issues concerning this communication, please contact Applicants' primary attorney-of record, Gregory A. Sebald (Reg. No. 33,280), at (612) 336.4728.

Respectfully submitted,

MERCHANT & GOULD P.C.
P.O. Box 2903
Minneapolis, Minnesota 55402-0903
(612) 332-5300

Dated: November 30, 2001

By 

Gregory A. Sebald
Reg. No. 33,280

GAS/tvm

MARKED-UP COPY

10. (Amended) A vector comprising the isolated DNA molecule as claimed in [any one of claims 1 to 9] claim 1, wherein said isolated DNA molecule is under control of a regulatory element that directs expression of said DNA in a plant cell.
15. (Amended) A transformed plant cell comprising the vector of [any one of claims 10 to 14] claim 10.
16. (Amended) A transformed plant comprising the vector of [any one of claims 10 to 14] claim 10.
18. (Amended) An isolated protein encoded by the isolated DNA molecule as claimed in [any one of claims 4 to 9] claim 4.
19. (Amended) A method of producing asexually derived embryos comprising:
- i) transforming a plant cell with the vector of [any one of claims 10 to 14] claim 10;
 - ii) growing said plant cell to produce transformed tissue;
 - iii) selecting said transformed tissue for occurrence of said isolated DNA molecule; and
 - iv) assaying said transformed plant for asexual embryo production.
25. (Amended) A method of modifying the regenerative capacity of a plant comprising
- i) transforming a plant cell with the vector of [any one of claims 10 to 14] claim 10;
 - ii) growing said transformed plant cell to produce transformed tissue; and
 - iii) assaying said transformed plant tissue for enhanced regeneration as compared to wild-type tissue.
27. (Amended) A method of selecting a transformed plant comprising;
- i) transforming a normally non-regenerative plant with a vector of [any one of claims 10 to 14] claim 10; and
 - ii) determining whether said transformed plant is able to regenerate under conditions in which said normally non-regenerative plant does not regenerate.

30. (Amended) A vector comprising the isolated DNA molecule of claim 28 [or 29] operably associated with a gene of interest, wherein said isolated DNA molecule directs the expression of said gene of interest within a plant cell.
33. (Amended) A transformed plant cell comprising the vector of [either] claim 30[, 31 or 32].
34. (Amended) A transformed plant comprising the vector of [either] claim 30[, 31 or 32].
36. (Amended) A method for directing the expression of a gene of interest within a developing embryo of a plant comprising transforming said plant with the vector as defined by [either] claim 30[, 31 or 32].
37. (Amended) A use of a nucleotide sequence as defined in [any one of claims 4, 5, 6 or 7] claim 4 as a selectable marker.
38. (Amended) A method of producing asexually derived embryos comprising:
- i) transiently transforming a plant cell with the vector of [any one of claims 10 to 14, or introducing into said plant cell the protein of claim 18] claim 10, to produce a modified plant cell;
 - ii) growing said modified plant cell to produce tissue; and
 - iii) assaying said tissue for asexual embryo formation.
44. (Amended) A method of modifying the regenerative capacity of a plant comprising
- i) transiently transforming a plant cell with the vector of [any one of claims 10 to 14, or introducing into said plant cell the protein of claim 18] claim 10, to produce a modified plant cell;
 - ii) growing said modified plant cell to produce tissue; and
 - iii) assaying said tissue for enhanced regeneration as compared to wild-type tissue.
46. (Amended) A method of producing an apomictic plant comprising:
- i) transforming a plant with the vector of [any one of claims 10 to 14] claim 10, to produce a transformed plant;

- ii) selecting said transformed plant for occurrence of said isolated DNA molecule; and
- iii) assaying said transformed plant for asexual embryo production.

51. (Amended) A method of modifying the regenerative capacity of a plant comprising

- i) transiently transforming a plant cell with the vector of [any one of claims 10 to 14, or introducing into said plant cell the protein of claim 18] claim 10;
- ii) growing said plant cell to form tissue; and
- iii) assaying said tissue for enhanced regeneration as compared to wild-type tissue.

53. (Amended) A method of selecting a modified plant comprising;

- i) transiently transforming a normally non-regenerative plant with a vector of [any one of claims 10 to 14, or introducing into said normally non-regenerative plant the protein of claim 18,] claim 10 to produce said modified plant; and
- ii) determining whether said modified plant is able to regenerate under conditions in which said normally non-regenerative plant does not germinate.


55. (Amended) A method of producing a protein of interest comprising

- i) transforming a plant with [at least on vector, said at least one vector selected from any one of claims 10 to 14] a vector of claim 10 to produce a transformed plant;
- ii) selecting said transformed plant for occurrence of said isolated DNA molecule; and
- iv) growing said transformed plant in order to produce said protein of interest, wherein expression of said protein of interest is induced by the expression product of said isolated DNA.

58. (Amended) The method of [any one of claims 55 or 56] claim 55, wherein said protein of interest is selected from the group consisting of a pharmaceutically active protein, antibody, industrial enzyme, protein supplement, nutraceutical, storage protein, an enzyme involved in oil biosynthesis, animal feed, and animal feed supplement.

59. (Amended) The isolated DNA molecule of [claim of any one of claims 4 to 7] claim 4, wherein said isolated DNA molecule encodes a protein that is at least 70% similar with the amino acid defined by SEQ ID NO:2.

60. (Amended) The isolated DNA molecule of [claim of any one of claims 4 to 7] claim 4, wherein said isolated DNA molecule encodes a protein that is at least 70% similar with the amino acid defined by SEQ ID NO:4.
- 66 (Amended) A vector comprising the isolated DNA molecule as claimed in [any one of claims 63 to 65] claim 63, wherein said isolated DNA molecule is under control of a regulatory element that directs expression of said DNA in a plant cell.
71. (Amended) A transformed plant cell comprising the vector of [any one of claims 66 to 70] claim 66.
72. (Amended) A transformed plant comprising the vector of [any one of claims 66 to 70] claim 66.



ABSTRACT

Title: USE OF THE BNM3 TRANSCRIPTIONAL ACTIVATOR TO CONTROL PLANT EMBRYOGENESIS AND REGENERATION PROCESS

The present invention provides for a gene obtained during the induction of microspore embryogenesis. The protein encoded by this gene renders plant cells embryogenic, and increases the regenerative capacity of the plant cell. Also disclosed is the regulatory region of this gene and its use for directing the expression of a gene of interest within a suitable host cell.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211

09/980364
Rec'd PCT/PTO 30 NOV 2001

USE OF THE *BNM3* TRANSCRIPTIONAL ACTIVATOR TO CONTROL PLANT
EMBRYOGENESIS AND REGENERATION PROCESSES

BACKGROUND OF THE INVENTION

A typical angiosperm seed consists of three major components, the embryo, the endosperm and the maternal seed coat. Seed development begins with a double fertilization event, in which one sperm cell nucleus fuses with the egg cell nucleus to form the embryo, and a second sperm cell nucleus fuses with two central cell nuclei to form the endosperm. Embryo development itself can be separated into three developmental phases. The first phase of embryo development is one of cell division and morphogenesis, which serves to establish the major tissue types and organ systems of the mature plant. The second phase encompasses a period of rapid cell expansion and is characterized by the synthesis of storage reserves that sustain the embryo during germination and early seedling development. In the final phase of embryo development, the embryo becomes desiccated and enters into a period of developmental arrest or dormancy. All of the above events normally take place while the seed remains attached to the maternal plant.

Many plant species are capable of producing embryos in the absence of fertilization. This process of asexual embryo development may occur naturally, for example on the leaf margins of *Bryophyllum* (Yarborough, 1923) and *Malaxis* (Taylor, 1967), or within the ovule of apomictic plants (Koltunow, 1995). Apomixis refers to the production of a seed from the maternal ovule tissues in the absence of egg cell fertilization. Asexual embryo development may also be induced *in vitro* from gametophytic or somatic tissue (Mordhorst *et al.*, 1997) or, as shown recently, may be induced by genetic modification of gene expression (Ogas *et al.*, 1997; Lotan *et al.*, 1998).

Three major mechanisms of apomixis, diplospory, apospory and adventitious embryony, have been observed. Each mechanism differs with respect to the source of the cell that gives rise to the embryo and with respect to the time during ovule development at which the apomictic process is initiated. Diplospory and apospory are considered gametophytic forms of apomixis as

they involve the formation of diploid embryo sacs. Adventitious embryony does not involve the production of a mitotically-derived embryo sac.

In diplospory, the megaspore mother cell does not undergo normal meiosis, but rather divides mitotically to produce a diploid embryo sac instead of the normal haploid embryo sac. One of the cells of the embryo sac functions as the egg cell and divides parthenogenetically (without fertilization) to form an embryo. In some species the unreduced polar nuclei of the embryo sac may fuse to form the endosperm (autonomous endosperm production), the nutritive tissue of the seed, while in other species pollination is necessary for endosperm production (pseudogamy).

In aposporous apomicts, parthenogenic embryos are produced from additional cells, the aposporous initials, that differentiate from the nucellus. As with the megagametophyte of diplosporous species, the aposporous initial undergoes mitotic divisions to produce a diploid embryo sac. Aposporous embryos are not derived from the megagametophyte and can therefore co-exist within a single ovule with sexually-derived embryos. Autonomous production of endosperm is rare in aposporous species. Aposporous apomicts therefore depend on fertilization of the polar nuclei of a meiotically-derived embryo sac for the production of endosperm.

With adventitious embryony, embryos are formed directly from sporophytic ovule tissue, such as the integuments or nucellus, via parthenogenesis. Seeds derived from species exhibiting adventitious embryony generally contain multiple asexually-derived embryos and may also contain a single sexually-derived embryo. Plants exhibiting adventitious embryo also rely on the presence of a meiotically-derived embryo sac within the same ovule for endosperm formation.

In most plant species, the apomictic trait appears to be under the control of a single dominant locus. This locus may encode one or more developmental regulators, such as transcription factors, that in sexually reproducing plants function to initiate gene expression cascades leading to embryo sac and/or embryogenesis, but which are heterochronically or ectopically expressed in apomictic plants (Peacock, 1995; Koltunow, 1993; Koltunow et al, 1995).

Apomixis is a valuable trait for crop improvement since apomictic seeds give rise to clonal offspring and can therefore be used to genetically fix hybrid lines. The production of hybrid seed is a labour intensive and costly procedure as it involves maintaining populations of genetically pure parental lines, the use of separate pollen donor and male-sterile lines, and line isolation. Production of seed through apomixis avoids these problems in that once a hybrid has been produced, it can be maintained clonally, thereby eliminating the need to maintain and cross separate parental lines. The use of apomictic seed also provides a more cost effective method of multiplying vegetatively-propagated crops, as it eliminates the use of cuttings or tissue culture techniques to propagate lines, reduces the spread of diseases which are easily transmitted through vegetatively-propagated tissues, and in many species reduces the size of the propagule leading to lower shipping and planting costs.

Although apomixis occurs in a wide range of plant species, few crop species are apomictic. Attempts to introduce apomictic traits into crop species by introgression from wild relatives (Ozias-Akins, *et al.*, 1993; WO 97/10704; WO 97/11167) or through crosses between related, but developmentally divergent sexual species (WO 98/33374), have not yielded marketable products. Other approaches have focused on the identification of gene sequences that may be used to identify or manipulate apomictic processes (WO 97/43427; WO 98/36090), however these approaches have not led to methods for the routine production of apomictic plants.

Mutagenesis approaches have also been attempted to convert sexually reproducing plants such as *Arabidopsis thaliana* (arabidopsis) into apomictic plants (Peacock *et al.*, 1995). For example, a number of recessive "fertilization-independent seed" (*fis*) mutants have been identified that initiate partial embryo and/or endosperm at a low frequency in the absence of fertilization (Chaudhury *et al.*, 1997). However, a number of additional parameters need to be modified in order to obtain true diploid apomictic seed using *fis* mutants.

Asexually-derived embryos can be induced to form in culture from many gametophytic and somatic plant tissues (Yeung, 1995). Somatic embryos can be obtained from culture of somatic tissues by treating them with plant growth regulators, such as auxins, or auxins in

combination with cytokinins. Embryos can also be induced to form in culture from the gametophytic tissues of the ovule (gynogenesis) and the anther (androgenesis, pollen or microspore embryogenesis), either by the addition of plant growth regulators or by a simple stress treatment.

Several mutants have been identified that may be used to induce efficient production of embryos *in vitro*. These include recessive arabidopsis mutants with altered shoot meristems, for example *primordia timing (pt)*, *clavata (clv)1* and *clv3*, which were shown to enhance embryogenic callus formation when seedlings were germinated in the presence of auxin (Mordhorst *et al.*, 1998). The altered expression of two arabidopsis genes, *LEAFY COTYLEDON (LEC1)*; WO 98/37184, Lotan *et al.*, 1998) and *pickle*, have been shown to promote the production of somatic embryos in the absence of added growth regulators. The *LEC1* gene encodes a homologue of the HAP3 subunit of a CCAAT box-binding transcription factor (CBF). The *LEC1* gene controls many aspects of zygotic embryo development including desiccation tolerance and cotyledon identity. Ectopic over-expression of the *LEC1* gene in a *lec1* mutant background results in the production of 2 transgenic lines that occasionally form embryo-like structures on leaves. These embryo-like structures express genes, such as those encoding seed storage proteins and oil body proteins, which are normally preferentially expressed in developing embryos. Plants containing a recessive mutant *PICKLE* gene produce a thickened, primary root meristem. Mutant *pickle* roots produce embryo-forming callus when the root tissue is separated from the rest of the plant and placed on minimal medium without growth regulators (Ogas *et al.*, 1997). Mutant *pickle* roots show morphological characteristics of developing seeds, such as oil bodies and, as with *LEC1* over-expressers, accumulate genes preferentially expressed in developing seeds.

Efficient production of apomictic seed is only likely to be realised through the identification and subsequent modification of developmental regulators, such as transcription factors, that are known to activate gene expression cascades leading to embryogenesis in both sexually-reproducing and apomictic plants. The present invention addresses this need by providing methods for the production of apomictic seeds comprising ectopic over-expression of an embryo-expressed AP2 domain containing transcription factor, BNM3 or its homologs.

- 5 -

It is an object of the invention to overcome disadvantages of the prior art.

The above object is met by the combinations of features of the main claims, the sub-claims disclose further advantageous embodiments of the invention.

SUMMARY OF THE INVENTION

The present invention relates to asexual embryo formation and regeneration in plants. More specifically, it relates to processes for producing asexually-derived embryos, and for enhancing regeneration capacity in plants.

According to the present invention there is provided an isolated DNA molecule comprising a nucleotide sequence that:

- hybridizes to SEQ ID NO:5 or 6, not including the AP2 repeat1-linker-AP2 repeat2 region, under moderate or stringent conditions;
- hybridizes to SEQ ID NO:1 or 3, not including the AP2 repeat1-linker-AP2 repeat2 region, under stringent conditions;
- comprises at least 27 contiguous nucleotides of SEQ ID NO's:1, 3, 5 or 6; or that
- exhibits at least 70% similarity with the nucleotide sequence defined by SEQ ID NO's:1, 3, 5 or 6.

This invention further relates to an isolated DNA molecule that hybridizes to SEQ ID NO's: 1, 3, 5 or 6, not including the AP2 repeat1-linker-AP2 repeat2 region, under moderate or stringent conditions, and comprises a nucleic acid sequence encoding a protein, wherein the protein when present at a sufficient level within a plant cell renders the cell embryogenic, increases the regenerative capacity of the plant cell, or both renders the cell embryogenic and increases the regenerative capacity of the plant cell. Included within the present invention is the isolated DNA molecule as just defined comprising a nucleotide sequence that hybridizes to nucleotides 1-2014 of SEQ ID NO:1, 1-2011 of SEQ ID NO:3, 1620-4873 of SEQ ID NO:5, or nucleotides 2026-5035 of SEQ ID NO:6, not including the AP2 repeat1-linker-AP2 repeat2 region, under moderate or stringent conditions. Also included within the present invention is a vector comprising the isolated DNA molecule as defined above, wherein the isolated DNA molecule is under control of a regulatory element that directs expression of said DNA in a plant cell. The regulatory element may be a constitutive, inducible, tissue specific or a developmental active, regulatory element.

This invention also embraces a transformed plant cell, a transformed plant, or seed obtained from a transformed plant, each comprising the vector as defined above

This invention relates to an isolated protein encoded by an isolated DNA molecule that hybridizes to the nucleotide sequence defined by SEQ ID NO:1, 3, 5 or 6, not including the AP2 repeat1-linker-AP2 repeat2 region, under moderate or stringent conditions, wherein the protein, when present at a sufficient level within a plant cell renders the cell embryogenic, or increases the regenerative capacity of the plant cell. Also included is a protein encoded by an isolated DNA molecule that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5, or nucleotides 2026-5035 of SEQ ID NO:6, not including the AP2 repeat1-linker-AP2 repeat2 region, under moderate or stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under moderate or stringent conditions. This invention also embraces an isolated DNA molecule that encodes a protein as defined by SEQ ID NO:2, SEQ ID NO:4 or SEQ ID NO:7. The invention also pertains to a protein comprising at least 70% homology with the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4 or SEQ ID NO:7, or comprises from about 30 to about 541 amino acids of the sequence disclosed in SEQ ID NO:2, or comprises from about 30 to about 561 amino acids of the sequence disclosed in SEQ ID NO:4.

The present invention is also directed to a method of producing asexually derived embryos comprising:

- i) transforming a plant cell with a vector comprising an isolated DNA molecule that hybridizes to SEQ ID NO's: 1, 3, 5, or 6, not including the AP2 repeat1-linker-AP2 repeat2 region, under moderate or stringent conditions and which encodes a protein that when present at a sufficient level within the plant cell renders the plant cell embryogenic, or increases the regenerative capacity of the plant cell, or that comprises a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5, or nucleotides 2026-5035 of SEQ ID NO:6, not including the AP2 repeat1-linker-AP2 repeat2 region, under moderate or stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under moderate or stringent conditions;
- ii) growing the plant cell to produce transformed tissue;

- iii) selecting the transformed tissue for occurrence of the isolated DNA molecule; and
- iv) assaying the transformed tissue for asexual embryo formation.

This invention also relates to the above method where the step of assaying (step iv)) involves assaying for somatic embryos, gametophytically-derived embryos, adventitious embryony, diplospory, or for haploid parthenogenesis of the embryo sac.

The present invention also embraces a method of producing an apomictic plant comprising:

- i) transforming a plant cell with a vector comprising an isolated DNA molecule that hybridizes to nucleotides 1-2014 of SEQ ID NO:1, nucleotides 1-2011 of SEQ ID NO:3, nucleotides 1620-4873 of SEQ ID NO:5, or nucleotides 2026-5035 of SEQ IDNO:6, not including the AP2 repeat1-linker-AP2 repeat2 region, under moderate or stringent conditions and which encodes a protein that when present at a sufficient level within said plant cell renders the plant cell embryogenic, or increases the regenerative capacity of the plant cell;
- ii) selecting the transformed plant for occurrence of the isolated DNA molecule; and
- iii) assaying the transformed plant for asexual embryo formation.

This invention also relates to the above method where the step of assaying (step iii)) involves assaying for asexually-derived embryos, somatic embryos, gametophytically-derived embryos, adventitious embryony, diplospory, or for haploid parthenogenesis of the embryo sac.

The present invention is also directed to a method of producing asexually derived embryos comprising:

- i) transiently transforming a plant cell with a vector comprising an isolated DNA molecule that hybridizes to nucleotides 1-2014 of SEQ ID NO:1, nucleotides 1-2011 of SEQ ID NO:3, nucleotides 1620-4873 of SEQ ID NO:5, or nucleotides 2026-5035 of SEQ IDNO:6, not including the AP2 repeat1-linker-AP2 repeat2 region, under moderate or under stringent conditions and which encodes a protein that when present at a sufficient level within the plant cell renders the plant cell embryogenic, or increases the regenerative capacity of the plant cell;

- ii) growing the transiently transformed plant cell to produce transiently transformed tissue;
- iii) assaying the transiently transformed tissue for asexual embryo formation.

This invention is directed to the above method where the step of assaying (step iii)) involves assaying for asexually-derived embryos, somatic embryos, gametophytically-derived embryos, adventitious embryony, diplospory, or for haploid parthenogenesis of the embryo sac.

The present invention also presents a method of modifying the regenerative capacity of a plant comprising

- i) transforming a plant cell with a vector comprising an isolated DNA molecule that hybridizes to SEQ ID NO:5 or SEQ ID NO:6 under moderate or stringent conditions and which encodes a protein that when present at a sufficient level within the plant cell renders the plant cell embryogenic, or increases the regenerative capacity of said plant cell, or that comprises a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5, or nucleotides 2026-5035 of SEQ ID NO:6, not including the AP2 repeat1-linker-AP2 repeat2 region, under moderate or under stringent conditions, or to the nucleotide sequence as defined within nucleotides 1-2014 of SEQ ID NO:1, or nucleotides 1-2011 of SEQ ID NO:3 under stringent conditions;
- ii) growing the transformed plant cell to produce transformed tissue; and
- iii) assaying the transformed tissue for enhanced regeneration as compared to wild type tissue.

This invention also embraces the above method wherein step iii) includes assaying in the absence of a growth regulator.

The present invention also relates to a method of modifying the regenerative capacity of a plant comprising:

- i) transiently transforming a plant cell with a vector comprising an isolated DNA molecule that hybridizes to SEQ ID NO:5 or SEQ ID NO:6 under moderate or stringent conditions and which encodes a protein that when present at a sufficient level within the plant cell renders the plant cell embryogenic, or increases the

regenerative capacity of the plant cell, or that comprises a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5, or nucleotides 2026-5035 of SEQ IDNO:6, not including the AP2 repeat1-linker-AP2 repeat2 region, under moderate or under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions;

- ii) growing the transiently transformed plant cell to produce transiently transformed tissue;
- iii) assaying the transformed tissue for enhanced regeneration as compared to wild type tissue.

This invention also embraces the above method wherein step iii) includes assaying in the absence of a growth regulator.

The present invention also relates to a method of selecting a transformed plant comprising;

- i) transforming a normally non-regenerative plant with a vector comprising an isolated DNA molecule that hybridizes to SEQ ID NO:5 or SEQ ID NO:6 under moderate stringent conditions and which encodes a protein that when present at a sufficient level within the plant cell renders the plant cell embryogenic, or increases the regenerative capacity of said plant cell, or that comprises a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5, or nucleotides 2026-5035 of SEQ IDNO:6, not including the AP2 repeat1-linker-AP2 repeat2 region, under moderate or under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions; and
- ii) determining whether the transformed plant is able to regenerate under conditions in which the normally non-regenerative plant does not regenerate.

The present invention is also directed to an isolated DNA molecule comprising a DNA sequence that exhibits at least about 70% similarity with nucleotides 1-1619 of SEQ ID NO:5, or nucleotides 1-2025 of SEQ ID NO:6, or that comprises at least 22 contiguous nucleotides within nucleotides 1-1619 of SEQ ID NO:5 or 1-2025 of SEQ ID NO:6. Also included within

the scope of the present invention is a vector comprising the isolated DNA molecule as just defined, operably associated with a gene of interest, wherein the isolated DNA molecule directs the expression of the gene of interest within a plant cell. The gene of interest may be heterologous with respect to the isolated DNA molecule. The gene of interest may be selected from the group consisting of a pharmaceutically active protein, antibody, industrial enzyme, protein supplement, nutraceutical, storage protein, animal feed and animal feed supplement. This invention also includes a transformed plant cell, a transformed plant, or seed obtained from the transformed plant, comprising the vector as just defined.

Furthermore, the present invention includes a method for directing the expression of a gene of interest within a developing embryo of a plant comprising transforming said plant with a vector containing an isolated DNA molecule that exhibits at least about 70% similarity with nucleotides 1-1619 of SEQ ID NO:5, or nucleotides 1-2025 of SEQ ID NO:6, or that comprises at least 22 contiguous nucleotides within nucleotides 1-1619 of SEQ ID NO:5 or nucleotides 1-2025 of SEQ ID NO:6.

This invention also pertains to a method of producing a protein of interest comprising

- i) transforming a plant with at least one vector, comprising an isolated DNA molecule that hybridizes to SEQ ID NO:5 or SEQ ID NO:6 under moderate or stringent conditions and which encodes a protein that when present at a sufficient level within the plant cell renders the plant cell embryogenic, or increases the regenerative capacity of said plant cell, or that comprises a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5, or nucleotides 2026-5035 of SEQ ID NO:6, not including the AP2 repeat1-linker-AP2 repeat2 region, under moderate or under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions to produce a transformed plant;
- ii) selecting the transformed plant for occurrence of the isolated DNA molecule; and
- iv) growing the transformed plant in order to produce the protein of interest, wherein expression of the protein of interest is induced by the expression product of said isolated DNA.

This method may also comprise transforming the plant with a second vector comprising a nucleotide sequence encoding the protein of interest under the control of a regulatory element, wherein the regulatory element induced by the expression product of the isolated DNA. Furthermore, this method may also be used to produce a protein of interest wherein the protein of interest is a native protein.

This summary of the invention does not necessarily describe all necessary features of the invention but that the invention may also reside in a sub-combination of the described features.

BRIEF DESCRIPTION OF THE DRAWINGS

These and other features of the invention will become more apparent from the following description in which reference is made to the appended drawings wherein:

Figure 1 shows a schematic representation of the effect of culture temperature on the developmental fate of isolated microspores and pollen of *Brassica napus*. Late uninucleate microspores and early binucleate pollen cultured at 25 °C or lower continue to divide and form functional pollen grains (gametophytic), while the same microspores and pollen cultured at 32 °C undergo numerous sporophytic divisions, leading to the formation of haploid embryos (embryogenic). Late uninucleate microspores and early binucleate pollen cultured for one day at 25 °C, followed by culture at 32 °C may undergo gametophytic divisions, but form neither embryos nor mature pollen grains (non-embryogenic).

Figure 2 shows the alignment of the DNA sequences depicted in SEQ ID NO:1 and SEQ ID NO:3. The ATG and TAG translation initiation and translation termination codons are shown in bold. Identical nucleotides are indicated by (*) and gaps are indicated by (-).

Figure 3 shows the alignment of the predicted protein sequences encoded by the DNA of SEQ ID NO:1 and SEQ ID NO:3. The amino acid sequence of the first AP2 domain repeat (repeat 1) and the second AP2 domain repeat (repeat 2), are shown in bold. Identical amino acids are indicated by an asterisk (*) and mismatches by a dot (.) below the sequence alignment.

Figure 4 shows the presence of two *BNM3* genes in the *Brassica napus* genome. A DNA gel blot containing restriction digests of *B. napus* c.v. Topas genomic DNA was hybridized to a *BNM3A* cDNA fragment under high stringency conditions. The *BNM3A* cDNA hybridizes to two DNA fragments under these conditions. These fragments correspond to the *BNM3A* and *BNM3B* genes. The position of the molecular size markers (Lambda

DNA *Hind* III restriction fragments) is indicated to the left the figure. The restriction enzymes used to digest the DNA are indicated above the blot.

Figure 5 shows the alignment of the predicted protein sequence encoded by the DNA of SEQ ID NO.1 (BNM3A) with the predicted protein sequences of other AP2 domain proteins. The amino acid sequence of BNM3A, beginning at position 208, and spanning the first AP2 domain repeat (AP2 domain repeat 1), the second AP2 domain repeat (AP2 domain repeat 2), and the linker region lying between the two repeats (linker), was aligned with the amino acid sequence of other proteins containing two AP2 domains. The amino acid similarity in this region ranges from 53% for APETALA2 to 80% for ZMMHCF1. Identical amino acids are indicated by (*) and gaps are indicated by (-). Protein names are indicated on the left and are abbreviated as follows: ANT, AINTEGUMENTA (accession number U41339); ZM, ZMMHCF1 (accession number Z47554); GL15, GLOSSY15 (accession number U41466); AP2, APETALA2 (accession number U12546).

Figure 6 shows the results of gel blot analysis with a *BNM3A* cDNA fragment performed on RNA extracted from the indicated tissues. RNA gel blots contain either 5 µg (a) or 20 µg (b, c) of total RNA. Figure 6A shows the pattern of *BNM3* expression in microspore embryo cultures. RNA was isolated from late uninucleate microspores and early binucleate pollen at the time of collection (pollen 0d), after four days in culture at 32 °C (+ embryo), after four days in culture at 25 °C (pollen 4d), after one day of culture at 25 °C, followed by three days of culture at 32 °C (- embryo) and microspore-derived embryos at the globular, heart, torpedo, 21 day old cotyledon (21 d cot), 28 day old cotyledon (28 d cot) and 42 day old cotyledon (42 d cot) stage of development. *BNM3* expression is detected in embryogenic microspores and developing microspore-derived embryos, but is absent from developing microspores and pollen collected prior to tissue culture and in non-embryogenic samples. The exposure time was seven days. Figure 6B shows that *BNM3* gene expression is detected in developing seeds. Seeds were collected at various days after pollination (DAP). These points in development correspond approximately to the globular (7 d), heart (14 d), torpedo (18 d), early cotyledon (21 d),

mid cotyledon (28 d, 35 d) and late cotyledon (42 d) stages of development. The exposure time was 14 days. Figure 6C shows that *BNM3* gene expression is not detected in non-seed tissues. Roots and leaves were collected from 14 day old greenhouse grown plants. Entire flowers as well as excised anthers and pistils were collected from opened flower buds just prior to anthesis. Small and large buds refer to closed flower buds of less than 5 mm or greater than 5 mm in length, respectively. Siliques were collected 16 days after pollination. The exposure time was 14 days.

Figure 7 shows the phenotype of *Brassica napus* and arabidopsis plants transformed with constructs containing the *BNM3* gene under control of a modified *POLYUBIQUITIN* promoter (B) and double enhanced *35S* promoter containing an AMV translational enhancer (A, C-E). Figure 7A shows embryo structures on the leaf margin of a *Brassica* T1 seedling. Figure 7B shows embryo structures on the petiole of an arabidopsis T2 seedling. Figure 7C shows embryo structures on the cotyledon of an arabidopsis T1 seedling. Figure 7D shows a scanning electron micrograph of the abaxial side of an arabidopsis T1 cotyledon. Note the bipolar nature of the embryos, as well as the emergence of a secondary embryo from the surface of a primary embryo (asterisk). Figure 7E shows a semi-thin section through one of the cotyledons of the T1 seedling shown in (Figure 7C). Note the presence of all the major organs and tissue elements of embryo, as well as the development of new embryos on the flanks of the shoot apical meristems and the cotyledons.

Figure 8 shows the increased regenerative capacity of arabidopsis plants transformed with a construct containing the *BNM3B* gene under control of a modified *POLYUBIQUITIN* promoter. Figure 8A shows wild-type and transgenic leaf and hypocotyl explants on medium containing growth regulators. Figure 8B shows wild-type and transgenic roots on medium containing growth regulators. Figure 8C shows wild-type and transgenic leaf and hypocotyl explants on medium without growth regulators. Figure 8D shows wild-type and transgenic root explants on medium without growth regulators.

Figure 9 shows characterization of the *AtBBM* gene; the arabidopsis orthologue of *BNM3*.

Figure 9A shows the position of the *AtBBM* sequence and selected restriction endonuclease sites on approximately 8 kb of overlapping *Arabidopsis thaliana* ecotype C24 genomic DNA. The predicted protein coding region of the single arabidopsis *BNM3* homologue spans positions 3426 to 6435 of the genomic sequence. The exons of the predicted coding region are shown as black boxes above the restriction map. A vertical arrow at position 7479 indicates the start of the *LXR3* (*Irregular xylem3*) sequence. The horizontal scale bar is in kilobases. Figure 9 B shows that the arabidopsis *AtBBM* genomic clones and the arabidopsis homologue identified through DNA gel blot analysis using a *Brassica napus* *BNM3* cDNA probe are the same. Genomic DNA from arabidopsis ecotypes Landsberg *erecta* (L) and Columbia (C) was digested with the restricted enzymes shown in (A) and hybridised to a full-length *BNM3A* cDNA probe (SEQ ID NO1). Comparison of the restriction map shown in (A) with the pattern of hybridising restriction fragments indicates that the full-length *Brassica* cDNA probe detects a single *Arabidopsis* homologue under moderate stringency wash conditions (0.2X SSC, 0.1% SDS at 25 °C). The molecular size marker (in kilobases) is indicated to the right of the blot.

DESCRIPTION OF PREFERRED EMBODIMENT

The present invention relates to asexual embryo formation and regeneration in plants. More specifically, it relates to processes for producing asexually-derived embryos, and for enhancing regeneration capacity in plants. The present invention also relates to heterologous protein production systems in plants, and the uses thereof.

The following description is of a preferred embodiment by way of example only and without limitation to the combination of features necessary for carrying the invention into effect.

Genes preferentially expressed during the induction of *Brassica napus* c.v. Topas microspore embryogenesis were isolated via subtractive screening. Seven independent cDNA clones, comprising six unique DNA sequences were found to be differentially expressed between cDNA libraries prepared from embryogenic and non-embryogenic microspore cultures. Several of these *BNM* (for *Brassica napus* microspore embryo) clones, *BNM3A* (SEQ ID NO:1) and *BNM3B* (SEQ ID NO:3), were characterized as described herein. *BNM3A* and *BNM3B* encode the amino acid sequences disclosed in SEQ ID NO:2, and SEQ ID NO:4, respectively. The genomic sequence of *BNM3A* (SEQ ID NO:5), including the regulatory region (nucleotides 1-1619 of SEQ ID NO:5), was also obtained. The arabidopsis orthologue of the *BNM3* gene, called *AtBBM*, was also identified. The genomic sequence of *AtBBM* is depicted in SEQ ID NO:6 while the predicted amino acid sequence is shown in SEQ ID NO:7.

"Regeneration", as used herein, refers to a morphogenetic response that results in the production of new tissues, organs, embryos, whole plants or fragments of whole plants that are derived from a single cell, or a group of cells. Regeneration may proceed indirectly via a callus phase or directly, without an intervening callus phase. "Regenerative capacity" refers to the ability of a plant cell to undergo regeneration.

By "embryogenic cell", it is meant a cell that has completed the transition from either a somatic or a gametophytic cell to a state where no further applied stimuli are necessary to produce an embryo.

By "regulatory element" it is meant those that include developmentally regulated, tissue specific, inducible and constitutive regulatory elements. A regulatory element that is developmentally regulated, or controls the differential expression of a gene under its control, is activated within certain organs or tissues of an organ at specific times during the development of that organ or tissue. However, some regulatory elements that are developmentally regulated may preferentially be active within certain organs or tissues at specific developmental stages, they may also be active in a developmentally regulated manner, or at a basal level in other organs or tissues within the plant as well, such regulatory elements are considered "tissue specific". Regulatory elements may be found either upstream, within, downstream, or a combination thereof, of the coding region of a gene.

An inducible regulatory element is one that is capable of directly or indirectly activating transcription of one or more DNA sequences or genes in response to an inducer. In the absence of an inducer the DNA sequences or genes will not be transcribed. Typically the protein factor, that binds specifically to an inducible regulatory element to activate transcription, is present in an inactive form which is then directly or indirectly converted to the active form by the inducer. The inducer can be a chemical agent such as a protein, metabolite, growth regulator, herbicide or phenolic compound or a physiological stress imposed directly by heat, cold, salt, or toxic elements or indirectly through the action of a pathogen or disease agent such as a virus. A plant cell containing an inducible regulatory element may be exposed to an inducer by externally applying the inducer to the cell or plant such as by spraying, watering, heating or similar methods.

A constitutive regulatory element directs the expression of a gene throughout the various parts of a plant and continuously throughout plant development. Examples of known constitutive regulatory elements include promoters associated with the CaMV 35S

transcript. (Odell et al., 1985, *Nature*, **313**: 810-812), the rice actin 1 (Zhang et al, 1991, *Plant Cell*, **3**: 1155-1165) and triosephosphate isomerase 1 (Xu et al, 1994, *Plant Physiol.* **106**: 459-467) genes, the maize ubiquitin 1 gene (Cornejo et al, 1993, *Plant Mol. Biol.* **29**: 637-646), the *Arabidopsis* ubiquitin 1 and 6 genes (Holtorf et al, 1995, *Plant Mol. Biol.* **29**: 637-646), and the tobacco translational initiation factor 4A gene (Mandel et al, 1995 *Plant Mol. Biol.* **29**: 995-1004).

By "gene of interest" it is meant any gene that is to be expressed in a transformed plant. Such a gene of interest may include, but is not limited to, a gene that encodes a pharmaceutically active protein, for example growth factors, growth regulators, antibodies, antigens, their derivatives useful for immunization or vaccination and the like. Such proteins include, but are not limited to, interleukins, insulin, G-CSF, GM-CSF, hPG-CSF, M-CSF or combinations thereof, interferons, for example, interferon- α , interferon- β , interferon- τ , blood clotting factors, for example, Factor VIII, Factor IX, or tPA or combinations thereof. A gene of interest may also encode an industrial enzyme, protein supplement, nutraceutical, or a value-added product for feed, food, or both feed and food use. Examples of such proteins include, but are not limited to proteases, oxidases, phytases chitinases, invertases, lipases, cellulases, xylanases, enzymes involved in oil biosynthesis etc. Other protein supplements, nutraceuticals, or a value-added products include native or modified seed storage proteins and the like.

The present invention is further directed to a chimeric gene construct containing a DNA of interest operatively linked to a regulatory element of the present invention. Any exogenous gene, or gene of interest, can be used and manipulated according to the present invention to result in the expression of the exogenous gene.

The activation of the expression of a gene of interest may also be under the control of a regulatory element that itself is activated by a BNM3 protein. For example, which is not to be considered limiting, a gene of interest may be fused to the napin promoter, and the napin promoter may be induced by BNM3. Furthermore, a gene of interest may be expressed within somatic tissues under the control of one or more

regulatory elements induced by BNM3, so that, as will be described in more detail below, the somatic tissue develops into a seed-like structure comprising embryogenic cells, and these seed-like structures produce the products of the gene of interest.

The chimeric gene construct of the present invention can further comprise a 3' untranslated region. A 3' untranslated region refers to that portion of a gene comprising a DNA segment that contains a polyadenylation signal and any other regulatory signals capable of effecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by effecting the addition of polyadenylic acid tracks to the 3' end of the mRNA precursor. Polyadenylation signals are commonly recognized by the presence of homology to the canonical form 5' AATAAA-3' although variations are not uncommon. Examples of suitable 3' regions are the 3' transcribed non-translated regions containing a polyadenylation signal of *Agrobacterium* tumor inducing (Ti) plasmid genes, such as the nopaline synthase (*Nos* gene) and plant genes such as the soybean storage protein genes and the small subunit of the ribulose-1, 5-bisphosphate carboxylase (ssRUBISCO) gene. The 3' untranslated region from the structural gene of the present construct can therefore be used to construct chimeric genes for expression in plants.

The chimeric gene construct of the present invention can also include further enhancers, either translation or transcription enhancers, as may be required. These enhancer regions are well known to persons skilled in the art, and can include the ATG initiation codon and adjacent sequences. The initiation codon must be in phase with the reading frame of the coding sequence to ensure translation of the entire sequence. The translation control signals and initiation codons can be from a variety of origins, both natural and synthetic. Translational initiation regions may be provided from the source of the transcriptional initiation region, or from the structural gene. The sequence can also be derived from the regulatory element selected to express the gene, and can be specifically modified so as to increase translation of the mRNA.

To aid in identification of transformed plant cells, the constructs of this invention may be further manipulated to include plant selectable markers. Useful selectable

markers include enzymes which provide for resistance to an antibiotic such as gentamycin, hygromycin, kanamycin, and the like. Similarly, enzymes providing for production of a compound identifiable by colour change such as *GUS* (β -glucuronidase), fluorescence, or luminescence, such as luciferase are useful.

Also considered part of this invention are transgenic plants containing a gene or chimeric gene construct of the present invention comprising a *BNM3* gene, a regulatory element obtained from *BNM3*, or the coding region from *BNM3* in operative association with a constitutive, developmental or inducible regulatory element, or a combination thereof. Methods of regenerating whole plants from plant cells are known in the art. In general, transformed plant cells are cultured in an appropriate medium, which may contain selective agents such as antibiotics, where selectable markers are used to facilitate identification of transformed plant cells. Once callus forms, shoot formation can be encouraged by employing the appropriate plant hormones in accordance with known methods and the shoots transferred to rooting medium for regeneration of plants. The plants may then be used to establish repetitive generations, either from seeds or using vegetative propagation techniques. The constructs of the present invention can be introduced into plant cells using Ti plasmids, Ri plasmids, plant virus vectors, direct DNA transformation, micro-injection, electroporation, biolistics etc. For reviews of such techniques see for example Weissbach and Weissbach, *Methods for Plant Molecular Biology*, Academy Press, New York VIII, pp. 421-463 (1988); Geierson and Corey, *Plant Molecular Biology*, 2d Ed. (1988); and Miki and Iyer, Fundamentals of Gene Transfer in Plants. In *Plant Metabolism*, 2d Ed. DT. Dennis, DH Turpin, DD Lefebvre, DB Layzell (eds), Addison Wesley, Langmans Ltd. London, pp. 561-579 (1997). The present invention further includes a suitable vector comprising the gene or the chimeric gene construct.

A class of genes have been isolated from *Brassica napus* microspore embryo cultures. These genes have been found to be important regulators of embryogenesis by their ability to induce the formation of asexually-derived embryos when ectopically expressed in the vegetative tissues of plants. These genes are hereinafter indicated as

BNM3 genes (*Brassica napus* microspore embryo). SEQ ID NO. 1 depicts the cDNA of *BNM3A*, SEQ ID NO. 3 depicts the cDNA of *BNM3B*, and the genomic sequence for *BNM3A* is given in SEQ ID NO:5. The regulatory region of *BNM3A* lies within nucleotides 1-1619 of SEQ ID NO:5. The predicted protein sequences encoded by the DNAs of SEQ ID NO. 1 and 3 are outlined in SEQ ID NOs. 2 and 4, respectively.

The orthologue of the *BNM3* gene has also been identified in arabidopsis and is hereinafter referred to as *AtBBM*. SEQ ID NO:6 depicts the genomic sequence for *AtBBM*. The regulatory region of *AtBBM* lies within nucleotides 1-2025 of SEQ ID NO:6. The predicted amino acid sequence is given in SEQ ID NO:7.

The *BNM3* translation products contain two copies of an AP2 domain separated by a linker region (Figure 3; amino acids 208-378 of SEQ ID NO: 2, and SEQ ID NO: 4, nucleotides 2694-4252 of SEQ ID NO:5, nucleotides 2938-4416 of SEQ ID NO:6), which herein is referred to as "AP2domain" or "AP2 domain repeat1-linker-AP2domain repeat2". The AP2 domain is thought to mediate protein-protein interactions. The ability of a number of AP2 domain containing proteins to bind DNA, coupled with the presence of putative nuclear localization signals and acidic regions that may function as transcriptional activators suggests these proteins function as transcription factors. The AP2 domain repeat1-linker-AP2domain repeat2 of *BNM3* exhibits about 99% homology with the AP2 domain of *AtBBM* (95% nucleotide similarity between *BNM3* and *AtBBM*), as well as a high degree of similarity with other AP2-comprising proteins, for example, ANT about 85% (76% nucleotide similarity), MOE17 (chromosome 3) about 85% (78% nucleotide similarity; if the introns are included in the comparison, for example the AP2 region from SEQ ID NO5, then MOE17 exhibits about 63% nucleotide similarity over a 285 bp region within the second AP2 domain), ZMMHCF1, 88%, or GLOSSY15, 66%. However, outside the two AP2 domains, the similarity of the sequence of *BNM3* and these other AP2-containing proteins decreases significantly.

By "*BNM3*" or "*BNM3* gene", it is meant the sequence of oligonucleotides as disclosed in SEQ ID NOs:1, 3, 5, or 6, or fragments, derivatives, or mutations thereof, or oligonucleotide sequences that exhibit at least:

- i) 70% homology or similarity, with a fragment or derivative of the sequences disclosed in SEQ ID NOs 1, 3, 5 or 6, not including the AP2 domain repeat1-linker-AP2 domain repeat2 region as defined by nucleotides 741-1257 of SEQ ID NO:1 nucleotides 672-1188 of SEQ ID NO:3, the corresponding sequence within coding region of nucleotides 2694-4252 of SEQ ID NO:5, or nucleotides 2938-4416 of SEQ ID NO:6 (the regions defined by nucleotides 2694-4252 of SEQ ID NO:5 or nucleotides 2938-4416 of SEQ ID NO:6, are interrupted by 7 introns; see Figure 9); or
- ii) 70% homology or similarity, with the full length of sequences disclosed in SEQ ID NOs 1, 3, 5 or 6 including the AP2 domain repeat1-linker-AP2 domain repeat2 region.

Such homology determinations may be made using oligonucleotide alignment algorithms for example, but not limited to a BLAST (GenBank URL: www.ncbi.nlm.nih.gov/cgi-bin/BLAST/), using default parameters: Program: blastn; Database: nr; Expect 10; filter: default; Alignment: pairwise; Query genetic Codes: Standard(1) or FASTA, again using default parameters. Using sequence similarity searches *AtBBM* exhibits about 85% homology with the full length of *BNM3*, and therefore, *AtBBM* is a *BNM3* gene. Furthermore, a *BNM3* gene may also be defined in terms of its ability to hybridize with sequences disclosed in the present invention. Therefore, "*BNM3*" or "*BNM3* gene", also includes:

- iii) oligonucleotides from greater than about 15 nucleotides in length, preferably 20 to 25 nucleotides in length, that associate with any of SEQ ID NO:1, 3, 5 or 6, or a fragment or derivative of the sequences disclosed in SEQ ID NOs 1, 3, 5 or 6, not including the region coding for the AP2 domain (AP2 domain repeat1-

linker-AP2 domain repeat 2) as defined above, under conditions of high stringency, for example, but not to be limited to, hybridization using gel blots (Southern hybridization) at about 65°C at 5X SSC, followed by wash conditions at 0.1X SSC, at 65°C; or

- iv) substantially full length nucleotide sequences, or nucleotide sequences of greater than about 1000 nucleotides in length, that associate with SEQ ID NO:1, or 3, or a fragment or derivative of the sequences disclosed in SEQ ID NOs 1, 3 of a length greater than about 1000 nucleotides, not including the region coding for the AP2 domain (AP2 domain repeat1-linker-AP2 domain repeat 2) as defined above, under conditions of moderate or high stringency, for example, but not to be limited to, hybridization using gel blots (Southern hybridization) at about 25°C at 0.2XSSC, 0.1% SDS, or 65°C 5XSSC, respectively, followed by wash conditions at 0.2XSSC, 0.1% SDS at 25°C.

Under conditions of moderate stringency, the full length BNM3 cDNA only hybridizes with fragments of *AtBBM* (see Figure 9B), and therefore, *AtBBM* is a BNM3 gene. Sequence analysis of the *AtBBM* genomic clones position the 5' end of the *IRREGULAR XYLEM3 (LXR3)* gene downstream of the putative *AtBBM* coding region (position 7479, Figure 9A). *LXR3* has previously been shown to map to a 150 kb region of chromosome 5 between the markers nga106 (33.26 cM) and mi438 (33.34 cM) (Taylor et al., 1999). This data indicates that the arabidopsis orthologue of the *Brassica BNM3* genes is encoded by a single gene that maps to chromosome 5. A sequence highly similar to *AtBBM*, TAMU BAC clone:T10B6 (accession number AP002073; Nakamura, May 18, 2000) has also been mapped to chromosome 5.

"*BNM3* gene" also includes DNA molecules that comprises at least 27 contiguous nucleotides of SEQ ID NOs:1, 3, 5, or 6 or at least 22 contiguous nucleotides within the regulatory region of nucleotides 1-1619 of SEQ ID NO:5, or nucleotides 1-2025 of SEQ ID NO:6. A fragment of *BNM3*, as defined may be used as a probe for the identification of nucleotides related to *BNM3* regulatory, or coding, regions within an organism, or as

primers for the amplification of these nucleotide sequences. Furthermore, molecules comprising at least 27 contiguous nucleotides, and preferably greater than about 30 to about 35 nucleotides, of the sequence of SEQ ID NOs: 1, 3, 5 or 6 and that encode a protein, or an active fragment thereof, that when present at a sufficient level within a plant cell renders the cell embryogenic, increases the regenerative capacity of the plant cell, or renders the cell embryogenic and increases the regenerative capacity of the plant cell, are also considered to be *BNM3* genes. Preferably, a *BNM3* gene comprises from about 50 to about 1981 nucleotides of SEQ ID NOs: 1 or 3, from about 50 to about 3538 nucleotides from the coding region (1620-4858) of SEQ ID NO:5, or from about 50 to about 3009 nucleotides from the coding region (2025-5035) of SEQ ID NO: 6.

The genomic *BNM3* sequences obtained from *Brassica napus* and arabidopsis are characterized as comprising 8 introns. These introns are found at nucleotides 1846-2298, 2720-2952, 3036-3160, 3170-3314, 3404-3553, 3628-3797, 3849-3961, and 4039-4148, of SEQ ID NO:5, and nucleotides 2249-2578, 2994-3220, 3304-3420, 3429-3521, 3611-3770, 3845-3969, 4020-4151 and 4229-4310 of SEQ ID NO:6. The start codon of SEQ ID NO's: 5 and 6 are at nucleotides 1620 and 2026, respectively, while the stop codons are found at positions 4856 and 5035, respectively.

By "*BNM3* regulatory region" it is meant the sequence of oligonucleotides that exhibit the property of regulating the expression of (either positively, for example an enhancer or promoter region, or negatively, for example a silencer region), and that are in operative association with, a *BNM3* gene. Typically the *BNM3* regulatory region comprises nucleotides upstream from the start site of a *BNM3* gene, however, sequences residing within other regions of the gene may also exhibit regulatory properties and be considered a *BNM3* regulatory region, for example but not limited to sequences within introns. An example of a *BNM3* regulatory region, which is not to be considered limiting, includes:

- a sequence operably linked with a *BNM3* gene that exhibits a regulatory function, for example but not limited to a regulatory region upstream from the start site of a *BNM3* gene or within an intron, or a fragment, derivative, or mutation thereof;
- nucleotides from about 1 to about 1619 in SEQ ID NO:5 or a fragment or derivative thereof;
- nucleotides from about 1 to about 2025 of SEQ ID NO:6 or a fragment or derivative thereof;
- a nucleotide sequence that associates with a nucleotide sequence from about 1000 to about 1619 of SEQ ID NO:5 or from about 1500 to about 2025 of SEQ ID NO:6, or a fragment or derivative thereof, under conditions of high stringency, for example, but not to be limited to, hybridization to gel blots at about 65°C in 5XSSC, followed by wash conditions at 0.1X SSC, 65°C;
- a nucleotide sequence that associates with a nucleotide sequence from about 1 to about 1000 of SEQ ID NO:5 or from about 1 to about 1500 of SEQ ID NO:6, or a fragment or derivative thereof, under conditions of moderate or high stringency, for example, but not to be limited to, hybridization to gel blots at about 25°C in 0.2XSSC, 0.1% SDS, or 65°C in 5XSSC, respectively, followed by wash conditions in 0.1X SSC, at 65°C; or
- a nucleotide sequence that exhibits at least 70% similarity with the nucleotide sequence from about 1000 to about 1619 of SEQ ID NO:5 or from about 1500 to about 2025 of SEQ ID NO:6, or a fragment thereof of at least about 22 nucleotides, as determined using oligonucleotide alignment (for example, but not limited to a BLAST or FASTA search, using default parameters; see above).

By "BNM3 protein" it is meant a protein, or a biologically active fragment thereof, that renders a plant cell embryogenic, increases the regenerative capacity of the plant cell, or renders the cell embryogenic, increases the regenerative capacity of the plant cell, and that is encoded by a *BNM3* gene, as defined above. Preferably, a *BNM3* protein comprises from about 30 to about 579 amino acids of the sequence disclosed in SEQ ID NO:2, from about 30 to about 579 amino acids of the sequence disclosed in SEQ

ID NO: 4, or from about 30 to about 581 amino acids of the sequence disclosed in SEQ ID NO:7. However, BNM3 protein may also be defined as a protein having at least 70% homology with either SEQ ID NO:2, 4, or 7 not including the AP2-repeat1-linker-AP2 repeat2 region (amino acids 208-378 of SEQ ID NO:s 2, and 3, amino acids 205-375 of SEQ ID NO:7).

Search of the sequence databases indicated that the *BNM3* translation products contain two copies of an AP2 domain (Figure 3; see also SEQ ID NO: 2 for BNM3A, SEQ ID NO: 4 for BNM3B, and SEQ ID NO:7 for AtBBM). The AP2 domain was first identified in APETALA2, an arabidopsis protein that regulates meristem identity, floral organ specification, seedcoat development and floral homeotic gene expression (Jofuku *et al.*, 1994), but has since been identified in a wide range of proteins with diverse functions.

The AP2 domain is usually between 58 to 68 amino acids in length and contains a conserved central core of 18 amino acids, characterized by its ability to form an amphipathic α helix, a structure thought to mediate protein-protein interactions. The ability of a number of AP2 domain containing proteins to bind DNA, coupled with the presence of putative nuclear localization signals and acidic regions that may function as transcriptional activators suggests these proteins function as transcription factors.

Two phylogenetically distinct classes of AP2 domain proteins have been identified; proteins with a single AP2 domain (EREBP-like) and proteins with two AP2 domains (AP2-like; (Zhou, 1997)). The proteins encoded by the genes of this invention represent unique members of the latter class of proteins.

Accordingly, an aspect of the present invention provides for an isolated DNA molecule that comprises a sequence encoding a protein that contains two AP2 domains. The protein, when present at a sufficient level in a plant cell, renders the cell embryogenic, increases the regenerative capacity of the cell, or both renders the cell embryogenic and increases the regenerative capacity of the cell.

Analysis of *BNM3* expression during microspore-derived embryo development, seed development, or non-seed tissue development, using Northern (Figure 6) indicated that the *BNM3* genes are preferentially expressed in embryogenic microspore cultures, microspore-derived embryos and seeds. *BNM3* transcripts were not detected in any of the non-seed tissues tested.

BNM3 mRNA is detected in microspore cultures induced to undergo embryogenesis, as well as in the subsequent globular, heart, torpedo and cotyledon stages of microspore-derived embryo development (e.g. Figure 6A). RNAs are also detected within developing seeds, 14 days after pollination (14 DAP), corresponding to the heart stage of embryo development. *BNM3* expression increases during the early (21 DAP) and mid-cotyledon (28 DAP) stages of embryo development and remains constant thereafter (Figure 6B).

Constitutive expression of *BNM3* resulted in the formation of somatic embryos on vegetative structures such as cotyledons, petioles, leaf blades and the shoot apical meristem of plants (Figure 7). In these experiments *BNM3* cDNAs were placed under the control of two separate constitutive promoter constructs, a modified sunflower *POLYUBIQUITIN* promoter construct, and a double enhanced 35S promoter construct containing an AMV translational enhancer, however, it is to be understood that any suitable constitutive promoter may be used for this purpose. Such *BNM3*-derived ectopic embryos contain all of the organ systems and tissue layers found in the developing zygotic embryo in that these embryos are bipolar (Figure 7E), consist of an axis, a hypocotyl and radicle region, shoot and root meristems, and cotyledons. In addition, each organ system contained the characteristic radial arrangement of three specialized tissue layers (epidermis, ground parenchyma and provascular tissue) found in zygotic embryos. Continued expression of the *BNM3* gene within the developing ectopic embryo leads to a reiteration of the embryo-forming process, with the result that new embryos are continuously formed on the surface of pre-existing embryos (Figure 7E).

Constitutive expression of *BNM3* results in the increased ability of a plant to regenerate shoots *in vitro* in the presence of added growth regulators. Root explants from transgenic plants ectopically expressing *BNM3* show at least a 5-fold increase in shoot regeneration in the presence of hormones as compared to root explants obtained from wild-type plants (Figure 8A,B). Shoots also developed faster in the transgenic explants, compared to the wild-type. Wild-type leaf and hypocotyl explants initially responded by producing callus on the cut end of the petiole (Figure 8B) followed by callus formation along the length of the petiole. In contrast, explants from transgenic lines immediately produced new shoots (Figure 8B) or roots from the cut end of the petiole. Explants that initially produced roots eventually also produced shoots.

Transgenic explants, constitutively expressing *BNM3* were also able to regenerate in the absence of added growth regulators. These explants, when placed on media lacking growth regulators regenerated shoots either from the cut end of the leaf and hypocotyl explants or from the nodule-like structures of root explants (Figures 8C,D). In all cases regenerated shoots developed, rooted, flowered and set seed. Conversely, wild-type leaf and hypocotyl explants placed on medium lacking growth regulators occasionally produce callus or roots at the cut end of the leaf petiole, however no shoots form from these structures (Figure 8C,D).

It is also considered within the scope of the present invention, that expression of *BNM3* may be used to initiate a developmental cascade within a transformed plant or plant cell. This cascade may arise as a result of the stable integration of a DNA-based vector expressing *BNM3* within a transformed plant, however, such a cascade may also arise as a result of transient expression of *BNM3*, and does not require the stable integration of the *BNM3*-based vector within a plant cell. These transient approaches may be useful for inducing somatic embryogenesis, gametophytically-derived embryogenesis, or increasing the regenerative capacity of a plant or plant cell.

Plants in which a *BNM3* gene is ectopically expressed exhibit advantageous qualities including:

- formation of asexually derived embryos;
- increased regenerative capacity of tissue explants;
- the ability of tissue explants to regenerate in the absence of added plant growth regulators; and
- the expression of seed components in non-seed organs in which *BNM3* is ectopically expressed.

Furthermore, plants that ectopically express at least one *BNM3* gene can be used for the production of recombinant proteins using seed specific regulatory elements.

For the applications of *BNM3* as described below, it will be advantageous to obtain a high level of the *BNM3* transcript and/or *BNM3* protein in order to obtain plants in which the phenotype is highly penetrant. This may be obtained by using genetic elements such as introns, transcriptional enhancers or translational enhancers which are known to enhance gene or protein expression levels.

The *BNM3* sequences of the present invention may be used for several applications including, but not limited to, the control of embryo processes, the control of regeneration processes, the use of regulatory sequences for targeted gene expression, the use of *BNM3* sequences as selectable markers of transformed plants, or for embryogenic cells. These applications are disclosed in more detail below.

Use of *BNM3* Sequences to Control Embryogenic Processes

As described herein, *BNM3* genes play an important role in initiation and maintenance of embryo development. *BNM3* genes have been found in a wide range of members of the plant kingdom. Regulatory regions obtained from these genes may be used to control the transcription of *BNM3* or a derivative or fragment thereof, or any gene of interest, using methods known to one of skill in the art.

Ectopic expression of a *BNM3* gene is sufficient to induce recurrent formation of asexually derived embryos on the vegetative tissues of plants (see example 4). Depending upon the promoter used, ectopic over-expression of *BNM3* genes may be used to produce somatic or gametophytic embryos. Somatic or gametophytic embryos may be obtained by expressing a *BNM3* gene under the control of a constitutive regulatory element, as is shown in Example 5, or may also be obtained by expressing a *BNM3* gene under the control of tissue specific or developmentally regulated elements, inducible elements derived from either plant or non-plant genes or through transient expression. In this respect, chemical induction systems (e.g. see Gatz and Lenk, 1998, which is incorporated by reference) or transient expression using methods which do not result in stable integration of the *BNM3* gene, or which make direct use of the *BNM3* protein e.g. microprojectile bombardment of DNA or protein may also be employed.

Temporal and/or spatial restriction of *BNM3* expression using inducible, tissue specific or developmentally regulated elements, is preferred when recurrent embryogenesis is not a desirable trait. The regulatory elements used to restrict *BNM3* to a specific developmental stage or cell type will depend on the application. For example, regulatory elements that may be used to express *BNM3* for the production of microspore-derived embryos include, but are not limited to, those of the class I low molecular weight heat shock inducible gene, *GMHSP17.3B* (Zarsky *et al.*, 1995, which is incorporated by reference), or microspore/pollen expressed genes such as *NTM19* (Custers *et al.*, 1997, EP 790,311, which are incorporated by reference), *BCP1* (Xu *et al.*, 1995, which is incorporated by reference), *LAT52* (Twell *et al.*, 1989, which is incorporated by reference), *BNM1* (Treacy *et al.* 1997, which is incorporated by reference) and *APG* (Roberts *et al.*, 1993, which is incorporated by reference).

Examples of regulatory elements that may be used to express *BNM3* for the production of somatic embryos include, but are not limited to, those of genes activated by plant growth regulators which are routinely used to induce somatic embryogenesis in tissue culture. Specific examples, which are to be considered non-limiting, include the cytokinin inducible *IB6* and *CKII* genes (Brandstatter and Kieber, 1998; Kakimoto,

1996, which are incorporated by reference) and the auxin inducible element, DR5 (Ulmasov *et al.*, 1997, which is incorporated by reference). However, it is to be understood that other regulatory elements may be included for the expression of *BNM3* in plants.

Furthermore, examples of gene regulatory elements suitable for directing expression of *BNM3* to obtain adventitious embryony include, but are not limited to, those obtained from the ovule and embryo expressed *SERK* gene (Schmidt *et al.*, 1997 which is incorporated by reference), the ovule expressed *AGL11* gene (Roundsley *et al.*, 1995, which is incorporated by reference), the nucellus expressed *NUC1* gene (Doan *et al.*, 1996; WO 98/08961, which are incorporated by reference), or the inner integument-expressed genes, *FBP7* (Angenent *et al.*, 1995, which is incorporated by reference) and *SC4* (US application 09/059,909, filed April 13, 1998, which is incorporated by reference) genes.

According to one aspect of the present invention there is provided a method for the efficient production of microspore-derived embryos in plants. This method involves:

- i) transforming a plant of interest, for example, *Brassica napus* (using transformation techniques known to one of skill, for example, DeBlock *et al.*, 1989, Clough and Bent 1998, Vergunst *et al.* 1998, Klein *et al.* 1987, which are incorporated herein by reference) with a vector construct, or isolated DNA, consisting of a *BNM3* gene under control of a suitable regulatory element, which may be constitutive, tissue specific, developmentally regulated, or inducible and, optionally, a marker gene for selection of transformants;
- ii) selecting transformed plants;
- iii) producing lines that ectopically overexpress the *BNM3* gene, or BNM3 protein;
- iv) isolating microspores and pollen from the transgenic lines and culturing microspores and pollen to induce embryogenesis.

Embryogenesis can be induced by any suitable protocol, for example, which is not to be considered limiting, culturing microspore and pollen for about four days at from about 28° to about 35 °C, preferably at about 32 °C, then transferring embryogenic cells or embryos to about 25 °C.

Using the above method, *Brassica napus* cultivars ectopically overexpressing *BNM3* show an increase in the percentage of embryogenic cells or embryos over that observed when microspores or pollen are prepared from wild-type plants that do not ectopically express *BNM3*.

Examples of regulatory elements that may be used to express *BNM3* for the production of microspore-derived embryos include, but are not limited to, those of the class I low molecular weight heat shock inducible gene, *GMHSP17.3B* (Zarsky *et al.*, 1995, which is incorporated by reference), or microspore/pollen expressed genes such as *NTM19* (Oldenhof *et al.*, 1996, EP 790,311, which are incorporated by reference), *BCP1* (Xu *et al.*, 1995, which is incorporated by reference), *LAT52* (Twell *et al.*, 1989, which is incorporated by reference), *BNM1* (Treacy *et al* 1997, which is incorporated by reference), and *APG* (Roberts *et al.*, 1993, which is incorporated by reference). Also useful are inducible regulatory elements, for example but not limited to, tetracycline-inducible promoter (Gatz 1997, which is incorporated by reference), steroid inducible promoter (Aoyama and Chua 1997, which is incorporated by reference) and ethanol-inducible promoter (Slater *et al* 1998, Caddick *et al.* 1998, which are incorporated by reference).

In a similar fashion, microspore-derived embryos may also be produced in plants by introducing into a plant of interest a *BNM3* protein, (e.g. via biolistics ; Klein *et al* 1987) and selecting for plants that exhibit increased microspore embryogenesis.

This invention also provides a method for the efficient production of somatic embryos *in vitro*. This method involves:

- i) transforming a plant, for example, arabidopsis using transformation techniques known to one of skill (for example, but not limited to, DeBlock *et al.*, 1989, Clough and Bent 1998, Vergunst *et al.* 1998, which are incorporated by reference), or a plant cell may also be transiently transformed using methods known to one of skill (for example, biolistics; Klein *et al.* 1987) with a vector construct containing a *BNM3* gene under control of suitable regulatory element, which may be constitutive, inducible or developmentally regulated, and, optionally, a marker gene for selection of transformants is transformed to several arabidopsis.
- ii) selecting transformed plants, and
- iii) culturing the desired explant from the selected transformed plants, for example, but not limited to, root, leaf or seedlings *in vitro*, in media with or without appropriate growth regulators, for example, but not limited to 2,4-D (e.g. Mordhorst *et al.*, 1998) to produce direct embryogenesis or embryogenic callus; and
- iv) transferring embryos, non-embryogenic callus, or both embryos and non-embryogenic callus to appropriate media for the production of embryos, plantlets, or both embryos or plantlets.

For example, when the results of the above method are compared with the production of somatic embryos *in vitro* using a number of arabidopsis ecotypes, directed embryogenesis or embryogenic callus is initiated at a higher frequency from transgenic lines ectopically over-expressing *BNM3* than in wild-type controls.

Examples of regulatory elements that may be used to express *BNM3* for the production of somatic embryos include, but are not limited to, those of genes activated by plant growth regulators which are routinely used to induce somatic embryogenesis in tissue culture. Specific examples, which are to be considered non-limiting, cytokinin inducible *IB6* and *CKII* genes (Brandstatter and Kieber, 1998; Kakimoto, 1996, which are incorporated by reference) and the auxin inducible element, DR5 (Ulmasov *et al.*, 1997, which is incorporated by reference). Also useful are inducible regulatory elements,

for example but not limited to, a tetracycline-inducible promoter (Gatz 1997, which is incorporated by reference), a steroid inducible promoter (Aoyama and Chua 1997, which is incorporated by reference), and an ethanol-inducible promoter (Slater et al 1998, Caddick et al. 1998, which are incorporated by reference).

Ectopic initiation of embryo development is one of the key steps in apomixis. As shown in Example 4, ectopic expression of a *BNM3* gene is sufficient to initiate embryo formation in otherwise non-embryo-forming tissue. A *BNM3* gene may therefore be used to initiate adventitious embryony or parthenogenesis of a reduced or unreduced embryo sac cell by expression of the gene in the sporophytic or gametophytic tissues of the developing ovule.

Adventitious embryony is achieved by expressing *BNM3* in sporophytic ovule tissues such as the nucellus, the inner integuments or other tissues lying adjacent to or in proximity to the developing embryo sac. This method involves:

- i) transforming a desired plant (see above methods) with a vector construct consisting of a *BNM3* gene under control of suitable regulatory element, which may be constitutive, inducible or developmentally regulated, and, optionally, a marker gene for selection of transformants, using methods known within the art;
- ii) selecting transformed plants;
- iii) emasculating the transformed plant;
- iv) pollinating the transformed plants with pollen carrying one or more dominant selectable markers, for example GUS or kanamycin resistance; and
- v) assaying for production of clonal offspring.

When the results of the above method are compared with the pollination of a wild-type arabidopsis plant with pollen carrying the dominant selectable marker, all F1 embryos resulting from this cross inherit the dominant marker while embryos derived

from plants ectopically over expressing the *BNM3* gene or protein are clonally derived via sexual embryo formation and do not inherit the dominant selectable marker.

Specific examples of gene regulatory elements suitable for directing expression of *BNM3* to obtain adventitious embryony, diplospory or haploid parthenogenesis of embryo sac components include the ovule expressed *SERK* gene (Schmidt et al. 1997, which is incorporated by reference), the meiosis expressed *AtDMC1* gene, (Klimyuk and Jones, 1997; WO 98/28431, which are incorporated by reference), the ovule expressed *AGL11* gene (Roundsley et al., 1995, which is incorporated by reference), the nucellus expressed *NUC1* gene (Doan et al., 1996; WO 98/08961, which are incorporated by reference), and the inner integument-expressed genes, *FBP7* (Angenent et al, 1995, which is incorporated by reference) and *SC4* (US application 09/059,909, filed April 13, 1998, which is incorporated by reference) genes. Furthermore, inducible systems, for example but not limited to, tetracycline-inducible promoter (Gatz 1997, which is incorporated by reference), steroid inducible promoter (Aoyama and Chua 1997, which is incorporated by reference), ethanol-inducible promoter (Slater et al 1998, Caddick et al. 1998, which are incorporated by reference) may also be used. Parthenogenesis from cells of the embryo sac requires a regulatory element that is active in one or more cells of the female gametophyte or their precursors. Fertilization of the meiotically-derived polar nuclei is desirable when the development of seed is dependent on the presence of endosperm.

Use of *BNM3* Sequences to Control Regeneration Processes

Plants ectopically over-expressing the *BNM3* genes exhibit increased regenerative capacity and the ability to regenerate whole plants in the absence of added growth regulators (see example 5). *BNM3* gene expression may therefore be used to enhance or induce the regeneration capacity of plant tissues *in vivo* or *in vitro*. The regulatory elements used to express *BNM3* will depend, in part, on the target tissue used for regeneration. Regeneration of plant tissues may be obtained by expressing a *BNM3* gene under the control of a constitutive regulatory element, for

example, but not limited to, 35S, or by expressing a *BNM3* gene under the control of tissue specific or developmentally regulated elements, inducible elements derived from either plant or non-plant genes (e.g. Gatz and Lenk, 1998, which is incorporated by reference), or through transient expression methods which do not result in stable integration of the *BNM3* gene or which make direct use of the *BNM3* protein (e.g. microprojectile bombardment of DNA or protein). Chemical induction systems (see Gatz and Lenk, 1998) or regulatory elements of genes that respond to plant growth regulators used to induce regeneration, such as, for example, cytokinin (Brandstatter and Kieber, 1998; Kakimoto, 1996) or auxin (Ulmasov et al., 1997), or genes expressed at the wound site of tissue explants (Xu et al., 1993) may be used.

A further application is the use of a *BNM3* gene as a selectable marker for the recovery of transgenic plants. As an example of this application which is not to be considered limiting in any manner, roots of a seedling, for example, an *Arabidopsis* ecotype C24 seedling, are cocultivated with a single *Agrobacterium tumefaciens* strain (per Vergunst et al, 1998; except that all steps are carried out in the absence of added growth regulators) containing two binary constructs:

- a first binary vector carries a reporter gene fusion, for example, but not limited to, 35S:*GUS*;
- a second binary vector contains a *BNM3* gene under control of suitable regulatory element.

BNM3 gene expression is activated upon integration of the above construct into the *Arabidopsis* genome and transgenic plants are selected on the basis of their ability to regenerate under conditions in which wild-type explants are unable to regenerate, for example, but not limited to, the absence of growth regulators. In many instances the T-DNA carrying the *BNM3* gene and the T-DNA carrying the gene of interest will integrate at unlinked loci. The T-DNA containing the introduced *BNM3* sequence, and its associated increased regenerative capacity phenotype, may therefore be removed in the progeny plants by simple segregation (Daley et al. 1998). However, as will be apparent to one of skill in the art, other methods such as transient

expression, which do not result in stable integration of the *BNM3* gene or which make direct use of the *BNM3* protein, may also be employed.

Use of *BNM3* Sequences to Target Gene Expression to the Embryo

Since *BNM3* genes are preferentially expressed in developing embryos (see example 3), a further application of this invention is the use of *BNM3* regulatory regions to target expression of at least one heterologous gene of interest to the developing embryo for any purpose, for example, but not limited to, altering embryo and seed traits such as seed viability or size, composition of constituents of the seed, disease resistance, or the production of high value products such as vaccines antibodies, biopharmaceuticals or other specialty chemicals.

Use of *BNM3* Expression as a Marker for Embryogenic Cells

As shown in Examples 3 and 4, *BNM3* gene expression is detected during the earliest phase of plant embryogenesis and is itself sufficient to activate signal transduction cascades leading to embryo development. *BNM3* gene expression is therefore a specific marker for the entry of a plant cell into the embryogenic pathway.

BNM3 expression is associated with embryo-forming cell divisions *in vitro* and *in vivo* and as such can be used to define culture conditions that alter the embryo-forming capacity of a tissue *in vitro*. Cells with embryogenic capacity or cells that undergo only a limited number of embryo-forming divisions are difficult to identify in the absence of structures that morphologically resemble embryos. However, these cells may be identified on the basis of *BNM3* expression. In this application, a vector containing the *BNM3* regulatory region, fused to a reporter gene, for example, but limited to, *GUS* (Jefferson *et al.*, 1987), *Luciferase* (Ow *et al.*, 1987) or *GFP* (Haselhoff and Amos, 1995) is transformed to a plant of interest. Homozygous transgenic lines exhibiting high levels of reporter gene expression in the embryo are cultured under *in vitro* conditions. Embryogenic cells, as well as culture conditions

which facilitate or enhance the formation of embryogenic cells are identified on the basis of reporter gene expression within the cultured tissue.

A related application is the use of the *BNM3* gene as a marker in apomictic species for the identification of individual cells that are in the process of forming asexually-derived embryos. In this application, cells entering the autonomous embryo pathway are identified by mRNA *in situ* hybridization using a RNA probe derived from a *BNM3* gene sequence, by immunocytochemistry using an antibody directed against a BNM3 protein, by transforming plants with a DNA construct containing a gene fusion between *BNM3* regulatory regions and a reporter gene, or by any similar technique known to those skilled in the art.

Identification of Signal Transduction Components

Signal transduction components which activate or are activated by *BNM3* gene expression can be elucidated by identifying proteins and DNA sequences that interact with a *BNM3* gene and its protein product. These signal transduction components may be identified using techniques known to a person skilled in the art, including for example, but not limited to:

- mutagenesis to identify intr- and/or extragenic suppressors or enhancers of the *BNM3* gain-of-function phenotype;
- yeast one hybrid screens for the isolation of proteins that bind to the *BNM3* regulatory regions to influence *BNM3* gene expression;
- genetic selection in yeast to identify genes that are direct targets of BNM3 binding;
- DNA arrays or proteomics to identify genes which are activated in a BNM3 signal transduction cascade; and
- yeast two hybrid screens to identify proteins that interact with BNM3 to influence expression of downstream target genes.

Techniques for the analysis of the signal transduction components and signalling components are well known (see for example, Meijer *et al.* (1998), Lipshutz *et al.* (1999), and Anderson and Anderson (1998)).

Plants over-expressing the *BNM3* gene under control of a strong constitutive regulatory element such as, for example, but not limited to, the Cauliflower Mosaic Virus 35S promoter exhibit ectopic embryo formation, enhanced regeneration via organogenesis or a combination thereof (Examples 4 and 5). The ability of *BNM3* ectopic over-expression to induce both embryo formation and enhance regeneration processes can be used to identify mutants altered in their embryo-forming or regenerative capacity. In this application a vector construct consisting of a *BNM3* protein coding region under control of a regulatory element that is sufficient to promote either ectopic embryo formation or enhanced regeneration phenotype is made and introduced into a plant of interest. Homozygous transgenic lines exhibiting a high penetrance of ectopic embryo formation, enhanced regeneration phenotype, or a combination thereof are identified. These lines are mutagenized by any available technique well known to the person skilled in the art, but which may include EMS mutagenesis, fast neutron mutagenesis, transposon mutagenesis or T-DNA mutagenesis. Mutagenized plants are then screened for alterations in the ectopic embryo formation or regeneration phenotype. These alterations include, for example, but not limited to, elimination or enhancement of the ability to promote ectopic asexual embryo formation or to regenerate in the absence of added growth regulators.

Heterologous Protein Expression System

Genetic control of the signal transduction pathway leading to embryogenesis and organogenesis in non-seed organs of transgenic plants may be activated by ectopic expression of a *BNM3* gene. Expression of a *BNM3* gene in association with a heterologous promoter can be used to produce altered seed components including for example, proteins, oils and other metabolites. Biotransformation of desired organs

may also include altering the nutritive value of, for example leaves of forage crops, or it may be used to create alternative uses for crops. The use of promoters that are induced by the signal transduction cascade initiated by expression of *BNM3* can be used to express high-valued recombinant proteins in organs other than seeds. An example of one such promoter is the napin promoter, obtained from the 2S seed storage protein napin. The production of proteins initiated from a *BNM3*-induced cascade, may be achieved within organs exhibiting greater biomass than seeds. Therefore, this technology may be used to create alternatives for plants as crops.

Accordingly, the present invention further relates to a binary system in which the *BNM3* protein binds directly or indirectly to an embryo-expressed regulatory sequence (target sequence) and activate transcription of a chimeric gene construct in any plant cell, tissue or organ. Therefore, *BNM3* may be used to directly or indirectly activate transcription of a chimeric gene construct. This approach involves *BNM3* interacting either directly with at least one target sequence from an embryo-expressed gene, or indirectly by initiating an embryogenic signal cascade that activates a transcription factor that in turn binds to and activates transcription from at least one target sequence. This binary system may be used for the expression of proteins in somatic tissues with the properties of expression in seeds.

In this application transgenic plants containing the *BNM3* gene under control of a constitutive regulatory element, for example, but not limited to the 35S promoter (35S:*BNM3*) are created to produce a *BNM3* activator line. *BNM3* expression may be demonstrated in a wide range of tissues in the *BNM3* activator lines by RNA gel blot analysis. Stable homozygous activator lines with high levels of *BNM3* expression are identified. Somatic tissues over-expressing *BNM3* may be examined for expression of other embryo-expressed genes, such as arabin (Guerche et al., 1990), cruciferin (Pang et al., 1988) or oleosin, or for morphological properties that are normally characteristic of seeds, such as the presence of lipid or protein bodies.

Transgenic plants of the same species to that used to generate the BNM3 activator lines described above are also created which contain an embryo-expressed promoter fused to a gene of interest, to produce a gene of interest line. In order to help describe this embodiment, the gene of interest line expresses a reporter gene, such as *GUS*, and examples, which are not to be considered limiting, of such lines include *Brassica napus* 2S albumin seed storage protein gene, *BngNAPI:GUS* fusion (Baszczynski *et al.*, 1994) or a *SERK:GUS* fusion (Schmidt *et al.*, 1997; a non-seed expressed reporter construct such as *BNM1:GUS* (Treacy *et al.*, 1997) may be used as a negative control). The fidelity of expression of the gene of interest in the specific organs and tissues of these gene of interest lines is demonstrated for each construct. Stable homozygous lines with high levels of expression of the gene of interest expression are created.

Transgenic lines containing BNM3 activator lines and gene of interest lines are crossed and the progeny seeds collected. *BNM3* gene expression, and in this example, *GUS* activity, expression of other embryo-expressed genes, as well as the morphological characteristics of transformed tissues, are examined. *BNM3* expression in non-seed tissues typically activates both embryo development and expression of the gene of interest (e.g. *GUS*), however, activation of the expression of the gene of interest in the absence of morphologically discernible embryos may also be observed. Expression of the gene of interest, in the absence of morphologically discernible embryos provides initial evidence for direct interaction of BNM3 with the target sequence.

Direct interaction of BNM3 with a target sequence may also be demonstrated using transient expression of BNM3 in plant protoplasts, along with the transient co-expression of an embryo-expressed promoter fused to a gene of interest (i.e. a gene of interest construct). *35S:BNM3* DNA and the gene of interest construct are introduced into protoplasts derived from non-seed cells, such as leaf mesophyll cells by electroporation. The expression of the gene of interest is examined after several hours to confirm activation of the target sequence. Direct interaction of BNM3 with the

target sequence may further be demonstrated by co-introducing the target sequence alone as competitor DNA.

In order to determine if tissues from different plant species may be transactivated by BNM3, 35S:BNM3 DNA and a reporter gene (for example, but not limited to *GUS*) construct may be introduced by microprojectile bombardment into somatic tissues of a plant. If BNM3 interacts directly with a target sequence then expression of the report er gene should coincide with transient expression of BNM3 in all species and tissues.

Direct evidence for BNM3-target sequence interaction may also be obtained by isolation of BNM3 protein expressed in bacteria, insect or yeast. *BNM3* is expressed in bacteria, insect, or yeast using commercially available expression systems and isolated to purity. Gel mobility shift assays (Gustavson *et al.*, 1991) are performed using a BNM3-target sequence, for example an embryo-expressed target sequence, to demonstrate direct binding of BNM3 to the BNM3-target sequence. Footprint analyses may also be performed to locate the region of BNM3 binding. Fragments of target sequences that bind BNM3 may then be subcloned and used as competitors for BNM3 binding in transient assays described above.

The following description is of a preferred embodiment by way of example only and without limitation to the combination of features necessary for carrying the invention into effect.

The present invention will be further illustrated in the following examples. However it is to be understood that these examples are for illustrative purposes only, and should not be used to limit the scope of the present invention in any manner.

Examples

General methods: *Microspore Embryo Culture*

Brassica napus c.v. Topas was used as the source of all plant material for microspore embryo culture. Donor plants for microspore culture were grown in a growth cabinet at 20 °C /15 °C (day/night) with a 16 h photoperiod (400 µE/m/s) provided by VHO cool white fluorescent lamps (165W, Sylvania) and incandescent bulbs (40W, Duro-test). Four weeks after germination the plants were transferred to growth cabinets under the same light conditions, but set at 10 °C /5 °C (day/night). Microspores and pollen were isolated and cultured as described in Keller *et al.* (1987), except that after 21 days in culture, cotyledon stage embryos were transferred to a maturation medium consisting of 1/2X NLN salts, 1% sucrose, 0.35 M mannitol and 5 µM ABA. Uninduced cultures (microspores and pollen continuing gametophytic development) and heat-stressed, non-embryogenic cultures (used for construction of the subtracted probe), were cultured from the same starting material as was used for the initiation of embryogenic cultures. Uninduced samples were obtained by culturing microspores and pollen for four days at 25 °C. Heat-stressed, non-embryogenic samples were obtained by culturing microspores and pollen for one day 25 °C, followed by three days 32 °C.

Samples of microspore and pollen cultured for less than 10 days were collected by centrifugation. Older samples containing globular, heart, torpedo and cotyledon stage microspore-derived embryos were collected by filtration through nylon meshes of various pore sizes as described in Ouellet *et al.* (1992). All other plant tissues were collected from greenhouse grown material. Seed material was obtained by hand pollinating flowers on the day of anthesis and collecting developing seeds on various days after pollination (DAP).

Nucleic Acid Isolation and Analysis

Total RNA was isolated using either a cesium chloride/guanidinium isothiocyanate procedure (Ouellet, 1992) or TRIZOL reagent (Gibco-BRL). RNA gel blot analysis was carried out by separation of 5 to 20 µg of total RNA per lane through 1.5% agarose gels containing 0.62 M formaldehyde, essentially as in Sambrook *et al.* (1989), followed by capillary transfer to Hybond-N nylon membranes (Amersham). Poly(A)⁺ RNA was isolated from total RNA by oligo (dT)-cellulose chromatography (Sambrook, 1989).

Genomic DNA was isolated from leaf tissue as described in Fobert *et al.* (1991) and digested with the specified restriction enzymes using standard procedures (Sambrook, 1989). DNA gel blot analysis was carried out by electrophoresis of 10 µg DNA through 0.8% agarose gels followed by capillary transfer to Hybond-N membranes.

The partial 1.2 kb *BNM3A* cDNA insert was used as a probe for DNA and RNA gel blots. Hybridization to gel blots was carried out at 65 °C according to the Hybond-N protocol. The final wash conditions were 0.1X SSC, 65 °.

Subtractive Probe Construction and cDNA Library Screening

Poly (A) mRNA was isolated from late uninucleate microspores and early binucleate pollen that had been cultured for four days at 32 °C in order to induce embryogenesis (embryogenic sample) and used to synthesize first strand cDNA (Riboclone cDNA kit; Promega). The cDNA was then hybridized to a five-fold excess (by weight) of poly (A)⁺ RNA from late uninucleate microspores and early binucleate pollen that had been cultured for one day at 25 °C, followed by three days at 32 °C to inactivate embryogenesis (non-embryogenic sample: Pechan *et al.*, 1991). The subtractive hybridization was performed essentially as described in Sambrook *et al.* (1989). The single-stranded cDNA recovered after subtraction was labelled with [α -³²P] dCTP using a random primers kit (BRL) and used as the subtracted probe for screening a Lambda phage cDNA library constructed from the same embryogenic

sample described above (Boutillier, 1994). Triplicate nylon filter lifts (Hybond-N) from approximately 1.5×10^5 plaque-forming units of the library were screened with the subtracted probe, with a random primers-labelled first strand non-embryogenic probe and with a random primers-labelled napin seed storage protein cDNA probe (pN2; Crouch, 1983). Napin mRNAs are prevalent in the embryogenic microspore library (Boutillier, 1994) and therefore plaques hybridizing to the napin probe were removed from the subsequent screening steps. Plaques hybridizing to the subtracted probe, but not to the non-embryogenic or napin probes, were selected and subjected to two subsequent rounds of differential screening using both the subtracted and non-embryogenic cDNA probes. DNA from selected Lambda clones was isolated (Sambrook, 1989), partially digested with *Eco* RI and *Xba* I and subcloned into pGEM-4Z (Promega).

Seven cDNAs comprising 6 unique genes, one of which comprised a truncated BNM3A cDNA, were identified. Two distinct, full length BNM3 cDNA clones (BNM3A and BNM3B) were subsequently obtained by stringent screening of circa 2.5×10^5 plaque-forming units of a cDNA library (UniZAPII cDNA synthesis kit, Stratagene) constructed with mRNA from 10 day old globular to heart-stage microspore-derived embryos of *B. napus* c.v. Topas. The BNM3 cDNA inserts were rescued by *in vivo* excision into Bluescript SK(-) (Stratagene).

Isolation of Brassica napus Genomic DNA sequences

The Universal Genome Walker Kit (Clontech) was used to isolate genomic DNA fragments lying upstream of the *BNM3* ATG start codon. Pools of uncloned, adaptor-ligated *Brassica napus* cv Topas genomic DNA fragments were constructed and used to isolate BNM3 genomic sequences by nested PCR. The primary PCR made use of the outer adaptor primer (AP1) supplied by the manufacturer and a BNM3 specific primer with the sequence:

5'-GAGGCAGCGGTCGGATCGTAACAGTACTCT-3' (SEQ ID NO:8).

The nested PCR made use of the nested adaptor primer (AP2) supplied by the manufacturer and a BNM3 specific primer with the sequence:

5'-CATAAGGAGAGAGAGAAAAGCCTAACCAGT - 3' (SEQ ID NO:9).

The primary PCR mixture was then diluted 1:50 and used as template for nested PCR. Both the primary and nested PCRs were performed as recommended by the manufacturer. The nested PCR products were cloned into the pGEMT-Easy vector (Promega) and sequenced. PCR products corresponding to the 5' untranslated genomic regions of both *BNM3A* and *BNM3B* cDNAs were identified.

The genomic DNA sequence spanning the *BNM3A* ATG translational start and TAG translational stop codons was isolated by PCR from *B. napus* cv Topas genomic DNA using *Pfu* polymerase (Stratagene) and the following primer combination:

5'-ACCAAGAACTCGTTAGATC-3' (SEQ ID NO:10); and

5'-AACGCATATAACTAAAGATC-3' (SEQ ID NO:11).

The primers were used under standard PCR conditions. The PCR products were cloned into the pGEMT-Easy vector and sequenced.

DNA Gel Blot Analysis and Mapping in Arabidopsis thaliana

Five hundred nanograms of arabidopsis genomic DNA (ecotypes Columbia and Landsberg *erecta*) (Shure et al., 1983) was digested with 20 different restriction endonucleases, separated by electrophoresis through 0.8% agarose gels and blotted onto Hybond N⁺ nylon membrane (Amersham) using standard methods. Blots were hybridised (1.5 M NaCl, 65 °C) with a ³²P[dATP] random primers labelled probe (Megaprime, Amersham) corresponding to either:

- 1) approximately the first 405 bp of the BNM3A cDNA (SEQ ID NO.1),
or

- 2) approximately the last 1200 nt of the *BNM3A* cDNA (SEQ ID NO.1) and then washed under conditions of low stringency (2 X SSC, 0.1% SDS at 65 °C) or moderate stringency (0.2 X SSC, 0.1% SDS at 25 °C).

A restriction fragment length polymorphism (RFLP) was identified between ecotypes Columbia and Landsberg *erecta* using the *Cfo* I restriction endonuclease. This RFLP was used to map the position of the arabidopsis BNM3 homologue on the arabidopsis genome using the Lister and Dean recombinant inbred (RI) lines (Lister and Dean, 1993). DNA from 100 recombinant inbred lines generated from a cross between ecotypes Columbia and Landsberg *erecta* was digested with *Cfo* I, transferred to Hybond N⁺ nylon membrane, hybridised with the *BNM3A* cDNA (as above) and washed under conditions of low stringency. The resulting RFLP data was sent to the RI database at the Nottingham Arabidopsis Stock Centre for determination of the map location (Lander et al., 1987). The results are discussed in Example 2-2, below.

Isolation of Arabidopsis thaliana Genomic DNA sequences

Three genome equivalents of an amplified arabidopsis ecotype C24 genomic Lambda phage library (Lambda-GEM 11, Promega) were screened using a truncated *BNM3A* cDNA probe (approximately the last 1200 nt of SEQ ID NO.1). Blots were hybridised with the ³²P-[dATP] random primers labelled probe (as above) and then washed under conditions of low stringency (2X SSC, 0.1% SDS at 65 °C). Seven Lambda phage were initially identified. Three putative full-length Lambda phage clones containing the arabidopsis homologue of the *BNM3* gene (SEQ ID NO:6) were subsequently identified after hybridisation under conditions of low stringency with a probe derived from the 5' end of the *Brassica napus* BNM3 cDNA (nt 1-405 of SEQ ID NO1). Individual clones comprising approximately 8.0 kb of overlapping sequence were identified from each of the three phage, subcloned into pBR322 and sequenced.

Plasmid Construction for Plant Transformation

The construction of a plasmid vectors containing the *BNM3* cDNAs under control of either a *POLYUBIQUITIN* or Cauliflower Mosaic virus 35S promoter are described below. The plasmid pRAP2TUBI contains a modified *Helianthus annuus* *POLYUBIQUITIN* promoter (Binet *et al.*, 1991) in the plasmid pRAP2T. The plasmid pRAP2T consists of the pUCAP plasmid (van Engelen *et al.*, 1995) and a nopaline synthase (nos) terminator inserted into the *Sac* I and *Eco* RI restriction sites. A PCR fragment of the *POLYUBIQUITIN* *Ubb1* promoter comprising the 5' end of the promoter to 7 bp from the 3' end of the first exon was amplified from the vector using an M13 reverse primer and the UBIQ-3' primer:

5'-CCATGGATCCAGAGACGAAGCGAAAC-3' (SEQ ID NO:12)

which includes introduced *Nco* I and *Bam* HI restriction sites. The *POLYUBIQUITIN* promoter fragment was digested with *Pst* I and *Bam* HI, gel purified and ligated into the *Pst* I and *Bam* HI sites of pRAP2T, creating the vector pRAP2TUBIH α . The full-length *BNM3B* cDNA was digested with *Eco* RI and *Xho* I restriction enzymes, blunted with Klenow enzyme, gel purified and ligated into the *Sma* I site of pRAP2TUBI making the plasmid pKB1S. An *Asc* I/*Pac* I DNA restriction fragment containing the modified *POLYUBIQUITIN* promoter, the *BNM3B* cDNA and the nos terminator was gel purified, and ligated to the *Asc* I/*Pac* I digested binary vector pBINPLUS (van Engelen *et al.*, 1995), creating the plasmid pKBBIN1S.

The construction of a vector containing the *BNM3A* cDNA under control of a double enhanced 35S promoter and AMV translational enhancer was as follows. A *Hind* III/*Xba* I DNA restriction fragment containing the double 35S promoter and the AMV translational enhancer from plasmid pBI525 (Datla *et al.*, 1993) was ligated to *Hind* III/*Xba* I digested pRAP2T, creating the plasmid pRAP2T35S. An *Nco* I site was introduced into the *BNM3A* cDNA clone by site directed mutagenesis. The sequence of the BNM3ANCO1 primer used for mutagenesis is:

5'-ACTCCATGGATAATAACTGGTTAGGC-3' (SEQ ID NO:13).

A second primer, BNM3AHINDIII:

5' - AAATTCTCAAGCTTTGGTCCATCTTG-3' (SEQ ID NO14)

was used together with the BNM3ANCO1 primer to amplify a 305 bp fragment of the *BNM3A* cDNA. This PCR fragment was digested with *Nco* I and *Hind* III and ligated to *Nco* I/*Kpn* I cut pRAP2T35S and a *Hind* III/*Kpn* I fragment containing the region of the *BNM3A* cDNA downstream of the *Hind* III site, creating the vector p35S:BNM3. p35S:BNM3 was digested *Asc* I and *Pac* I restriction enzymes and the fragment containing the double 35S promoter, the AMV translational enhancer, the *BNM3A* cDNA and the nos terminator was gel purified and ligated to the *Asc* I/*Pac* I digested binary vector pBINPLUS, creating the plasmid p35S:BNM3BIN.

Both the pKBBIN1S and p35S:BNM3BIN plasmids were transferred to *Agrobacterium tumefaciens* C58C1 strain carrying the disarmed Ti plasmid pMP90 and used in transformation experiments.

Plant Transformation

Arabidopsis thaliana ecotype C24 was used as the recipient in transformation experiments. Plants were transformed using either the floral dip method described in Clough and Bent (1998) or the root transformation method described in Vergunst *et al.* (1998).

Transgenic *Brassica napus* c.v. "Topas" plants were produced by *Agrobacterium tumaciens*-mediated transformation of microspore-derived embryos. Microspore-derived embryos were cultured for 5 weeks at a density of approximately 1000 embryos per ml. Overnight cultures of *Agrobacterium* were diluted 100 times in B5 medium containing 9% sucrose. Embryos were co-cultivated with the diluted bacteria for 48 hours at 24 °C in darkness, with slow shaking. The embryos were then transferred to NLN13 medium supplemented with 350 mg/L cefotaxim and 200 mg/L vancomycin for at least two weeks in darkness at 25 °C.

Embryos were germinated in weak light at 25 °C for about 2 weeks on solid B5 medium supplemented with 2% sucrose, cefotaxim (200 mg/L) and vancomycin

(100 mg/L). Well developed hypocotyls from germinated embryos were isolated and transferred to fresh germination medium supplemented with 100 mg/L kanamycin. After two weeks on this medium, explants were subcultured to a similar medium supplemented with kanamycin (25 mg/L). Green, putative transgenic, secondary embryos become visible after one month of selection.

Microscopy

All plant material was fixed overnight at 4 °C in 0.1 M phosphate buffer pH 7.0 containing 4% paraformaldehyde. Samples were washed in 0.1 M phosphate buffer and then dehydrated in a graded ethanol series to 100% ethanol. Samples for scanning electron microscopy were critical point dried in liquid CO₂ (Balzers CPD020), and mounted on SEM stubs using conductive carbon glue. Samples were coated with 30 nm palladium/gold using a Polaron E5100 sputter coater. Samples were observed in a JEOL JSM 5200 scanning electron microscope with an acceleration voltage of 15 kV. Digital images were obtained using Orion Framegrabber. Samples for light microscopy were embedded in Technovit 7100 (Kulzer). Sections were stained for 10 seconds in 1% Toluidine blue in 1% sodiumtetraborate, rinsed with water and mounted in Euparal. Digital images were recorded using a Sony 3 CCD camera.

Regeneration Experiments

Wild-type and transgenic arabidopsis seeds were surface sterilized, plated on ½ MS media containing 20% sucrose (½MS-20) and grown at 21 °C with the plates inclined at a 60° angle. Eight wild-type seedlings and eight seedlings from each of seven independent transgenic lines were harvested 10 days after germination and separated into root, hypocotyl and leaf explants. This material was then divided into two batches. Half of the explants were continuously cultured on B5 media containing 20% glucose (B5-20). Explants were transferred to fresh B5-20 media every two weeks. The remaining explants were cultured on B5-20 containing plant growth

regulators in order to induce shoot regeneration (Vergunst *et al.*, 1998). These explants were first placed on callus inducing media (CIM; high auxin to cytokinin ratio) for two days and then transferred to shoot inducing media (SIM; high cytokinin to auxin ratio) for the remainder of the culture period. Explants were transferred to fresh SIM media every two weeks.

Example 1: Isolation and Characterization of the BNM3 Genes from Brassica napus

A subtractive screening approach was used to isolate genes preferentially expressed during the induction of *Brassica napus* c.v. Topas microspore embryogenesis (Figure 1). Two types of microspore cultures were used in the construction of a subtracted probe: embryogenic and non-embryogenic. Embryogenic cultures were obtained by subjecting late uninucleate microspores and early binucleate pollen to a 4 day, 32 °C heat stress treatment. The non-embryogenic sample was obtained by culturing the same starting population of late uninucleate microspores and early binucleate pollen for 1 day at 25 °C followed by 3 days at 32 °C (Pechan *et al.*, 1991). Poly(A) mRNA was isolated from the embryogenic sample and used to synthesize first strand cDNA. The cDNA was then hybridized to an excess of poly(A)⁺ RNA isolated from a non-embryogenic microspore/pollen sample. The non-hybridizing, single stranded cDNA, enriched for sequences present in the embryogenic sample, but absent or present at a much lower level in the non-embryogenic sample, was recovered, radioactively labelled and used as a subtracted probe for screening a cDNA library derived from the embryogenic sample described above. Plaques hybridizing to the subtracted probe, but not to a probe derived from the non-embryogenic sample, were selected and subjected to two subsequent rounds of differential screening. Seven independent cDNA clones, comprising six unique DNA sequences were found to be differentially expressed between the embryogenic and non-embryogenic samples. One of these clones, 42A1, later renamed *BNM3A* (for *Brassica napus* microspore embryo), was further characterized.

Example 2-1: The BNM3 genes encode new members of the AP2 domain class of transcriptional activators

A single *BNM3* cDNA clone, *BNM3A*, was isolated after screening an embryogenic microspore cDNA library with a subtracted probe enriched for genes expressed in embryogenic microspores and pollen. The discrepancy between the size of the cDNA clone (1.2 kb) and the size of the transcript detected on RNA gel blots (2.2 kb) indicated that this clone did not represent a full-length cDNA. Two longer cDNA clones, corresponding to the full length cDNA of the clone originally isolated, *BNM3A* (SEQ ID NO. 1), and a new clone, *BNM3B* (SEQ ID NO. 3), were isolated from a 10 day old *Brassica napus* microspore embryo cDNA library. The alignment of the DNA sequence of these clones is shown in Figure 2. The two *BNM3* cDNA clones are 2011 and 1992 nt in length, and are 97% similar at the nucleotide level, differing only slightly in the length and sequence of their 5' and 3' untranslated regions. Both cDNAs potentially encode 579 amino acid polypeptides (predicted molecular mass of 63.9 kDa, pI of 5.7) that are 97% similar at the amino acid level (Figure 3).

The genomic complexity of the *BNM3* genes was determined by hybridization of the *BNM3* cDNAs to gel blots containing *B. napus* genomic DNA (Figure 4). The *BNM3* cDNAs hybridize to two DNA fragments under high stringency conditions. The two hybridizing fragments represent the two *BNM3* genes, *BNM3A* and *BNM3B*. *B. napus* is an amphidiploid species derived from the hybridization of the diploid *B. rapa* and *B. oleracea* genomes, thus the two *BNM3* sequences are likely derived from a single copy locus in each of the parental diploid progenitors.

Search of the sequence databases indicated that the *BNM3* translation products contain two copies of an AP2 domain (Figure 3). The AP2 domain was first identified in APETALA2 (AP2), an arabidopsis protein that regulates meristem identity, floral organ specification, seedcoat development and floral homeotic gene expression (Jofuku *et al.*, 1994; WO 98/07842), and has since been identified in a wide range of

proteins with diverse functions. These functions range from the activation of genes involved in stress (Zhou, 1997; Stockinger, 1997) and ethylene response (Ohme-Takagi, 1995) to the regulation of leaf, floral and ovule development (Moose, 1996; Jofuku, 1994; Elliot, 1996; Klucher, 1996). The AP2 domain is a 56-68 amino acid repeated motif containing at least two conserved regions: a highly basic YRG element, containing a conserved YRG amino acid motif and the RAYD element. The RAYD element contains a conserved central core of 18 amino acids that is predicted to form an amphipathic α -helix, a structure that is thought to mediate protein-protein interactions. The ability of a number of AP2 domain containing proteins to bind DNA, coupled with the presence of putative nuclear localization signals and acidic regions that may function as transcriptional activators suggests these proteins function as transcription factors.

Two phylogenetically distinct classes of AP2 domain proteins, consisting of either one AP2 domain (EREBP-like) or two AP2 domains connected by a linker region (AP2-like), have been identified (Zhou, 1997). BNM3 belongs to the latter class. Search of the databases with the region corresponding to the two AP2 domains and linker region of BNM3 reveals that BNM3 is most similar to the arabidopsis AINTEGUMENTA (ANT; Elliot, 1996; Klucher, 1996) and the *Zea mays* ZMMHCF1 AP2 domain containing protein. (ZM; Daniell, 1996) Figure 5 shows an alignment of the two AP2 domains of BNM3 with those of other proteins that contain two AP2 domains. BNM3 shares 85% amino acid sequence similarity with ANT and 88% with ZMMHCF1 in this region, but only 66% amino acid similarity with AP2 and GLOSSY15 in this region. A 10 amino acid insertion in the first AP2 domain of the BNM3 proteins further distinguishes these three proteins from other AP2 domain containing proteins (Elliot, 1996). The BNM3, AINTEGUMENTA and ZMMHCF1 proteins also share a small hydrophobic amino acid motif, LG/SFSL, in their amino terminal regions, but otherwise show no significant similarity in their DNA or amino acid sequences outside of the AP2 domains and linker. These results indicate that the BNM3 sequences encode unique members of the AP2 domain family of proteins.

A pairwise alignment of BNM3B cDNA and amino acid, sequences with ANT or ZMMHCF-1 sequences indicated that for the BNM3B nucleotide sequence:

- there is a 56% identity with ANT cDNA (over the 1905 nucleotides of ANT) and a 58% identity with ZMMHCF1 cDNA (over the 1773 nucleotide sequence of ZM);

and for the BNM3B amino acid sequence:

- there is a 41% identity of the BNM3B protein with ANT protein (over the 555 amino acid sequence of ANT), and a 46% identity with ZMMHCF1 protein (over 485 amino acid sequence of ZM).

Example 2-2: The Brassica napus BNM3 genes are represented by a single Arabidopsis thaliana orthologue

DNA gel blot analysis of arabidopsis genomic DNA hybridised to a number of *Brassica napus BNM3A* cDNA (SEQ ID NO:1) probes under conditions of low and moderate stringency indicated the presence of a single homologue of the *Brassica napus BNM3* genes in the arabidopsis genome. An RFLP was also identified between ecotypes Columbia and Landsberg *erecta* using the *Cfo* I restriction endonuclease. This RFLP was used to map the position of the single *BNM3* homologue on the arabidopsis genome to approximately 34 cM on chromosome 5 (Lister and Dean 1993).

Screening of three genomic equivalents of an arabidopsis genomic library identified three Lambda clones containing the putative full length arabidopsis *BNM3* homologue (*AtBBM*). Sequence analysis of the three *AtBBM* clones indicated that they are identical (SEQ ID NO:6). Figures 9A and B show respectively the restriction fragment pattern of the isolated arabidopsis genomic clones and the pattern of restriction fragments obtained after hybridisation of arabidopsis genomic DNA with the *Brassica BNM3A* cDNA probe. Comparison of the two figures indicates that the arabidopsis *AtBBM* genomic clones and the homologue identified through DNA gel blot analysis using a heterologous probe are the same. Sequence analysis of the

AtBBM genomic clones also positioned the 5' end of the *IRREGULAR XYLEM3* (*LXR3*) gene downstream of the putative *AtBBM* coding region (position 7479, Figure 9A). *LXR3* has previously been shown to map to a 150 kb region of chromosome 5 between the markers nga106 (33.26 cM) and mi438 (33.34 cM; Taylor et al., 1999). Together this data indicates that the arabidopsis orthologue of the *Brassica BNM3* gene is encoded by a single gene that maps to chromosome 5. A sequence that is very similar with *AtBBM*, TAMU BAC clone:T10B6 (accession number AP002073; Nakamura, May 18, 2000) also maps to chromosome 5.

Comparison of the structure of *Brassica BNM3A* genomic clone (SEQ ID NO:5) and the arabidopsis *AtBBM* genomic clone (SEQ ID NO:6) indicate that the predicted intron/exon boundaries are highly conserved between the two sequences. Both sequences are predicted to comprise nine exon and eight intron sequences.

Comparison of the DNA sequence of the *AtBBM* gene (SEQ ID NO:6) and the two *Brassica* cDNA sequences (SEQ ID NO:1 and SEQ ID NO:3) indicated that the three sequences are 85% similar across the entire putative protein coding region and 95% similar in the 546 nt region spanning the two AP2 domains and the linker region lying between the two AP2 domains. The nucleotide similarity to other related AP2 domain encoding genes such as *ANT* and the sequence located on clone MOE17 on chromosome 3 (accession number AB025629) in the region spanning the two AP2 domains and the linker region lying between the two AP2 domains is 76% and 78% respectively. Neither *ANT* nor the sequence on chromosome 3 shows significant DNA similarity to *AtBBM* outside of the AP2 domain encoding region.

The similarity between the predicted amino acid coding sequence of *AtBBM* and the two *Brassica* cDNA sequences (SEQ ID NO:1 and SEQ ID NO:3) is approximately 80% across the entire protein coding sequence and approximately 99% in the 182 amino acid region spanning the two AP2 domains and the linker region lying between the two AP2 domains. The amino acid similarity to both *ANT* and the AP2 domain sequence located on clone MOE17 on chromosome 3 in the amino acid

region spanning the two AP2 domains and the linker region lying between the two AP2 domains is approximately 85%. Neither of these two protein sequences shows significant similarity to AtBBM outside of the AP2 domain region of the protein.

Example 3: The BNM3 genes are preferentially expressed in developing embryos

RNA gel blot analysis (Figure 6) was used to determine the pattern of *BNM3* gene expression during microspore-derived embryo development, seed development, and in non-seed tissues. Both analyses indicate that the *BNM3* genes are preferentially expressed in developing embryos.

RNA gel blot analysis indicates that *BNM3* mRNAs are detected in microspore cultures induced to undergo embryogenesis, as well as in the subsequent globular, heart, torpedo and cotyledon stages of microspore-derived embryo development (Figure 6A). *BNM3* mRNAs are not detected in non-embryogenic microspore cultures, in freshly isolated microspores and pollen, or in microspores and pollen continuing gametophytic development in culture (Figure 6A). RNA gel blot analysis of developing seeds shows that *BNM3* expression is first detected 14 days after pollination (14 DAP), corresponding to the heart stage of embryo development. *BNM3* expression increases during the early (21 DAP) and mid-cotyledon (28 DAP) stages of embryo development and remains constant thereafter (Figure 6B). *BNM3* transcripts were not detected in any of the non-seed tissues tested, reflecting the low level or absence of transcripts in these tissues.

Example 4: Expression of BNM3 in Vegetative Tissues Promotes Asexual Embryo Formation

In order to determine the function the *Brassica napus* *BNM3* proteins, the *BNM3* cDNAs were placed under the control of two separate constitutive promoter constructs, a modified sunflower *POLYUBIQUITIN* promoter construct (hereafter referred to as *UBI:BNM3*) and a double enhanced 35S promoter construct containing

an AMV translational enhancer (hereafter referred to as 35S:*BNM3*), and introduced into arabidopsis. Analysis of the phenotype of the transformants indicates that ectopic over expression of the *BNM3* cDNAs promotes the formation of somatic embryos on vegetative structures such as cotyledons, petioles, leaf blades and the shoot apical meristem (Figure 7). The frequency of transformants producing ectopic embryos, as well as the penetrance of the ectopic embryo phenotype, was greater when the *BNM3* gene was expressed under control of the stronger double enhanced 35S promoter-AMV translational enhancer, as compared to the *POLYUBIQUITIN* promoter. Thus a high threshold level of protein product is required to increase the frequency and penetrance of the ectopic embryo phenotype.

BNM3-derived ectopic embryos contain all of the organ systems and tissue layers found in the developing zygotic embryo. *BNM3*-derived ectopic embryos are bipolar (Figures 7D and E) and consist of an axis, comprised of the hypocotyl and radicle regions, shoot and root meristems, and cotyledons (Figure 7E). In addition, each organ system contains the characteristic radial arrangement of three specialized tissue layers (epidermis, ground parenchyma and provascular tissue) found in zygotic embryos (Figure 7E). Continued expression of the *BNM3* gene within the developing ectopic embryo leads to a reiteration of the embryo-forming process, with the result that new embryos are continuously formed on the surface of pre-existing embryos (Figure 7D and E). These results provide conclusive evidence that expression of a single gene, *BNM3*, is sufficient to initiate a signal transduction cascade leading to the formation of fully differentiated asexually-derived embryos.

Example 5: Expression of BNM3 Increases the Regeneration Capacity of Plant Tissues

We examined the effect of *BNM3* gene expression on the ability of arabidopsis plants to regenerate shoots *in vitro* in the presence or absence of added growth regulators. Leaf, root and hypocotyl explants from 10 day old seedlings of wild-type arabidopsis and transgenic arabidopsis lines expressing *BNM3* under control of the

POLYUBIQUITIN promoter were placed on media containing growth regulators to induce first callus formation and then shoot organogenesis. Root explants from transgenic lines show at least a 5-fold increase in shoot regeneration in the presence of hormones as compared to wild-type root explants. (Figure 8A). These shoots also developed faster in the transgenic explants as compared to the wild-type. Wild-type leaf and hypocotyl explants responded by producing callus on the cut end of the petiole (Figure 8B). In contrast, explants from transgenic lines immediately produced new shoots (Figure 8B) or roots from the cut end of the petiole. Transgenic explants that initially produced roots eventually also produced shoots.

Transgenic explants were also able to regenerate in absence of added growth regulators. Wild-type leaf and hypocotyl explants placed on medium lacking growth regulators occasionally produced callus or roots at the cut end of the leaf petiole, however shoots did not regenerate from these structures (Figure 8C,D). Wild-type roots greened and formed thickened nodule-like structures at the junction with lateral roots, but did not develop further. In contrast, transgenic explants placed on media lacking growth regulators regenerated shoots either from the cut end of the leaf and hypocotyl explants or from the nodule-like structures of root explants (Figure 8C,D).

All citations are herein incorporated by reference.

The present invention has been described with regard to preferred embodiments. However, it will be obvious to persons skilled in the art that a number of variations and modifications can be made without departing from the scope of the invention as described herein.

References

Anderson, N.L. and Anderson, N.G. (1998). Proteome and proteomics: new technologies, new concepts and new words. *Electrophoresis* 19, 1853-1861.

Angenent, G.C., Franken, J., Busscher, M., van Dijken, A., van Went, J.L., Dons, H.J.M. and van Tunen, A.J. (1995). A novel class of MADS box genes involved in ovule development in *Petunia*. *Plant Cell* 7, 1569-1582.

Aoyama, T. and Chua, N.H. (1997). A glucocorticoid-mediated transcriptional induction system in transgenic plants. *Plant J.* 2, 397-404.

Baszczynski, C.L., and Fallis, L. (1990). Isolation and nucleotide sequence of a genomic clone encoding a new *Brassica napus* napin gene. *Plant Mol. Biol.* 14, 633-635.

Binet, M.N., Lepetit, M., Weil, J.H. and Tessier, L.H. (1991). Analysis of a sunflower polyubiquitin promoter by transient expression. *Plant Sci.* 79, 87-94.

Boutillier, K.A., Gines, M.J., Demoor, J.M., Huang, B., Baszczynski, C.L., Iyer, V.N. and Miki, B.L. (1994). Expression of the BnmNAP subfamily of napin genes coincides with the induction of *Brassica* microspore embryogenesis. *Plant Mol. Biol.* 26, 1711-1723.

Brandstatter, I. and Kieber, J.J. (1998). Two genes with similarity to bacterial response regulators are rapidly and specifically induced by cytokinin in *Arabidopsis*. *Plant Cell* 10, 1009-1019.

Caddick, M.X., Greenland, A.J., Jepson, I., Krause, K.P., Qu, N., Riddell, K.V., Salter, M.G., Schuch, W., Sonnewald, U., and Tomsett, A.B. (1998). An ethanolic inducible gene switch for plants used to manipulate carbon metabolism. *Nature Biotech.* 16, 177-180.

Chaudury, A.M., Letham, D.S., Craig, S. and Dennis, E.S. (1993). *amp-1-a* mutant with higher cytokinin levels and altered embryonic pattern, faster vegetative growth, constitutive photomorphogenesis and precocious flowering. *Plant J.* 4, 907-916.

- Chaudhury, A., Ming, L., Miller, C., Craig, S., Dennis, E. S., and Peacock, W. J. (1997). Fertilization-independent seed development in *Arabidopsis thaliana*. Proc. Natl. Acad. Sci. USA 94, 4223-4228.
- Clough, S.J. and Bent, A.F. (1998). Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. Plant J. 16, 735-743.
- Custers, J.B.M., Oldenhof, M.T., Schrauwen, J.A.M., Cordewener, J.H.G., Wullems, G.J. and van Lookeren Campagne, M.M. (1997) Analysis of microspore-specific promoters in tobacco. Plant Mol. Biol 35, 689-699.
- Crouch, M.L., Tenbarge, K.M., Simon, A.E. and Ferl, R. (1983). cDNA clones for *Brassica napus* seed storage proteins: evidence from nucleotide sequence analysis that both subunits of napin are cleaved from a precursor polypeptide. J. Mol. Appl. Genet. 2, 273-283.
- Daley, M., Knauf, V.C., Summerfelt, K.R. and Turner, J.C. (1998). Co-transformation with one *Agrobacterium tumefaciens* strain containing two binary plasmids as a method for producing marker free transgenic plants. Plant Cell Rep. 17, 489-496.
- Daniell, T.J., Fordham-Skelton, A.P., Vergani, P. and Edwards, R. (1996). Isolation of a maize cDNA (accession no. Z47554) (PGR 96-013) encoding APETALA-2-like binding domains by complementation cloning of an L-isoaspartyl methyltransferase-deficient mutant of *Escherichia coli*. Plant Phys. 110, 1435.
- Datla, R.S.S., Bekkaoui, F., Hammerlindl, J.K., Pilate, G., Dunstan, D.I. and Crosby, W.L. (1993). Improved high-level constitutive foreign gene expression in plants using an AMV RNA4 untranslated leader sequence. Plant Sci. 94, 139-149.

DeBlock, M. DeBrower, D. and Tenning, P. (1989). Transformation of *Brassica napus* and *Brassica oleracea* using *Agrobacterium tumefaciens* and the expression of the bar and neo genes in the transgenic plants. *Plant Physiol.* 91: 694-701.

Dellaert, L.M.W. (1981) Comparison of X-ray and fast neutron-induced mutant spectra. Experiments in *Arabidopsis thaliana* (L.) Heynh. *Arabidopsis Inf. Ser.*, 18, 16-36.

Doan, D. N. P., Linnestad, C., and Olsen, O.-A. (1996). Isolation of molecular markers from the barley endosperm coenocyte and the surrounding nucellus cell layers. *Plant Mol. Biol.* 31: 877-886.

Elliot, R.C., Betzner, A.S., Huttner, E., Oakes, M.P., Tucker, W.Q.J., Gerentes, D., Perez, P. and Smyth, D.R. (1996). AINTEGUMENTA, an APETALA2-like gene of *Arabidopsis* with pleiotropic roles in ovule development and floral organ growth. *Plant Cell* 8, 155-168.

Federoff, N., Furtak, D., and Nelson O. (1984). Cloning of the bronze locus in maize by a simple and general procedure using the transposable controlling element Ac. *Proc. Natl. Acad. Sci. USA* 81, 3825-3829.

Feldman, K.A., Marks, M.D., Christianson, M.L., and Quatrano, R.S. (1989). A dwarf mutant *Arabidopsis* generated by T-DNA insertion mutagenesis. *Science* 243, 1351-1354.

Fobert, P.R., Miki, B.L., and Iyer, V.N. (1991). Detection of gene regulatory signals in plants revealed by T-DNA-mediated fusions. *Plant Mol. Biol.* 17, 837-851.

Gatz, C. (1997). Chemical control of gene expression. *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 48, 89-108.

Gatz, C. and Lenk, I.R.P. (1998). Promoters that respond to chemical inducers. Trends Plant Sci. 3, 352-358.

Guerche, P., Tire, C., Grossi De Sa, F., De Clercq, A., Van Montagu, M. and Krebbers, E. (1990). Differential expression of the Arabidopsis 2S albumin genes and the effect of increasing gene family size. Plant Cell 2, 469-478.

Gustavsson, H.O., Ellerstrom, M., Stulberg, K., Ezcurra, I., Koman, A., Hoglund, A., Rask, L. and Josefsson, L.-G. (1991). Distinct sequence elements in a napin promoter interact in vitro with DNA-binding proteins from Brassica napus. Physiol. Plant 82, 205-212.

Haseloff, J. and Amos, B. (1995). GFP in plants. Trends Genet. 11, 328-329.

Jefferson, R.A. and Bicknell, R. (1996). The potential impacts of apomixis: a molecular genetics approach. In The Impact of Plant Molecular Genetics, B.W.S. Sobral, ed (Boston: Birkhauser), pp. 87-101.

Jefferson, R.A., Kavanagh, T.A. and Bevan, M.W. (1987). GUS fusions: β -glucuronidase as a sensitive and versatile gene fusion marker in higher plants. EMBO J. 6, 3901-3907.

Jofuku, K.D., den Boer, B.G.W., van Montagu, M. and Okamoto, J.K. (1994). Control of Arabidopsis flower and seed development by the homeotic gene APETALA2. Plant Cell 6, 1211-1225.

Kakimoto, T. (1996). CKI7, a histidine kinase homolog implicated in cytokinin signal transduction. Science 274, 982-985.

Keller, W.A., Fan, Z., Pechan, P., Long, N., Grainger, J. (1987). An efficient method for culture of isolated microspores of *Brassica napus*. Proceedings of the 7th International Rapeseed Congress. Poznan, Poland. Vol. 1, 152-157.

Klein, T.M., Wolf, E.D., Wu, R. and Sanford, J.C. (1987). High-velocity microprojectiles for delivering nucleic acids into living cells. *Nature* 327,70-73

Klimyuk, V.I. and Jones, J.D.G. (1997). AtDMC1, the Arabidopsis homologue of the yeast DMC1 gene: characterization, transposon-induced allelic variation and meiosis-associated expression. *Plant J.* 11, 1-14.

Klucher, K.M., Chow, H., Reiser, L. and Fischer, R.L. (1996). The AINTEGUMENTA gene of Arabidopsis required for ovule and female gametophyte development is related to the floral homeotic gene APETALA2. *Plant Cell* 8, 137-153.

Koltunow, A.M., Bicknell, R.A. and Chaudhury, A.M. (1995). Apomixis: molecular strategies for the generation of genetically identical seeds without fertilization. *Plant Physiol.* 108,1345-1352.

Koltunow, A.M. (1993). Apomixis: embryo sacs and embryos formed without meiosis or fertilization in ovules. *Plant Cell* 5, 1425-1437.

Korneef, M., Hanhart, C.J. and Thiel, F. (1989). A genetic and phenotypic description of eceriferum (cer) mutants in *Arabidopsis thaliana*. *J. Hered.* 80, 118-122.

Lander, E. S., Green, P., Abrahamson, J., Barlow, A., Day, M. J., Lincoln, S. E., and Newber, L. (1987). Mapmaker: an interactive computer package for constructing primary genetic linkage maps of experimental and natural populations. *Genetics* 121, 174-181.

- Lightner J., and Caspar, T. (1988) Seed Mutagenesis of Arabidopsis. In Arabidopsis Protocols, J.M. Martinez-Zapater and J. Salinas eds (Totowa, USA: Humana Press).
- Lipshultz, R.J., Fodor, S.P.A., Gingeras, T.R. and Lockhart, D.J. (1999). High density synthetic oligonucleotide arrays. *Nature Genetics* 21, 20-24.
- Lister, C., and Dean, C. (1993). Recombinant inbred lines for mapping RFLP and phenotypic markers in Arabidopsis thaliana. *Plant Journal* 4, 745-750.
- Lotan, T., Ohto, M., Matsudaira Yee, K., West, M.A.L., Lo, R., Kwong, R.W., Yamagishi, K., Fischer, R.L., Goldberg, R.B. and Harada, J.J. (1998). Arabidopsis LEAFY COTYLEDON1 is sufficient to induce embryo development in vegetative cells. *Cell* 93, 1195-1205.
- Meijer, A.H., Ouwerkerk, P.B.F. and Hoge, J.H.C. (1998). Vectors for transcription factor cloning and target site identification by means of genetic selection in yeast. *Yeast* 14, 1407-1415.
- Moose, S.P. and Sisco, P.H. (1996). Glossy15, an APETALA2-like gene from maize that regulates leaf epidermal cell identity. *Genes Dev.* 10, 3018-3027.
- Mordhorst, A.P., Toonen, M.A.J. and de Vries, S.C. (1997). Plant embryogenesis. *Crit. Rev. Plant Sci.* 16, 535-576.
- Mordhorst, A.P., Voerman, K.J., Hartog, M.V., Meijer, E.A., van Went, J., Koomneef, M., and de Vries, S.C. (1998). Somatic embryogenesis in Arabidopsis thaliana is facilitated by mutations in genes repressing meristematic cell divisions. *Genetics* 149, 549-563.

- Ogas, J., Cheng, J.-C., Sung, R.Z. and Somerville, C. (1997). Cellular differentiation regulated by gibberellin in the *Arabidopsis thaliana* pickle mutant. *Science* 277, 91-94.
- Ohme-Takagi, M. and Shinshi, H. (1995). Ethylene-inducible DNA binding proteins that interact with an ethylene responsive element. *Plant Cell* 7, 173-182.
- Oldenhof, M.T., de Groot, P.F.M., Visser, J.H., Schrauwen, J.A.M. and Wullems, G.J. (1996). Isolation and characterization of a microspore-specific gene from tobacco. *Plant Mol. Biol.* 31, 213-225.
- Ouellet, T., Rutledge, R.G. and Miki, B.L. (1992). Members of the acetohydroxyacid synthase multigene family of *Brassica napus* have divergent patterns of expression. *Plant J.* 2, 321-330.
- Ow, D.W., Jacobs, J.D. and Howell, S.H. (1987). Functional regions of the cauliflower mosaic virus 35S RNA promoter determined by use of the firefly luciferase gene as a reporter for promoter activity. *Proc. Natl. Acad. Sci. USA* 84, 4870-4874.
- Ozias-Akins, P., Lubbers, E.L., Hanna, W.W. and McNay, J.W. (1993). Transmission of the apomictic mode of reproduction in *Pennisetum*: co-inheritance of the trait and molecular markers. *Theor. Appl. Genet.* 85, 632-638.
- Pang, P., Pruitt, R. and Meyerowitz, E. (1988). Molecular cloning, genomic organisation, expression and evolution of the 12S storage protein genes of *Arabidopsis thaliana*. *Plant Mol. Biol.* 11, 805-820.
- Peacock, W.J., Ming, L., Craig, S., Dennis, E., Chaudury, A.M. (1995). A mutagenesis programme for apomixis genes in *Arabidopsis*. In *Proceedings Symposium on Induced Mutations and Molecular Techniques for Crop Improvement*, (Vienna: International Atomic Energy Agency), pp, 117-125

Pechan, P.M., Bartels, D., Brown, D.C.W. and Schell, J. (1991). Messenger-RNA and protein changes associated with induction of Brassica microspore embryogenesis.

Planta 184, 161-165.

Roberts, M.R., Foster, G.D., Blundell, R.P., Robinson, S.W., Kumar, A., Draper, J. and Scott, R. (1993). Gametophytic and sporophytic expression of an anther-specific *Arabidopsis thaliana* gene. Plant J. 3, 111-120.

Rounsley, S.D., Ditta, G.S. and Yanofsky, M.F. (1995). Diverse roles for MADS box genes in *Arabidopsis* development. Plant Cell 7, 1259-1269.

Salter, M.G., Paine, J.A., Riddell, K.V., Jepson, I., Greenland, A.J., Caddick, M.X., Tomsett, A.B. (1998). Characterisation of the ethanol-inducible alc gene expression system for transgenic plants. Plant Journal 16, 127-132.

Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989). Molecular Cloning: A Laboratory Manual, second edition. (Cold Spring Harbor: Cold Spring Harbor Laboratory Press).

Schmidt, E.D.L., Guzzo, F., Toonen, M.A.J. and de Vries, S.C. (1997). A leucine-rich repeat containing receptor-like kinase marks somatic plant cells competent to form embryos. Development. 124, 2049-2062.

Shure, M., Wessler, S., and Fedoroff, N. Molecular identification and isolation of the Waxy locus in maize. Cell. 1983 Nov; 35, 225-33.

Stockinger, E.J., Gilmour, S.J. and Thomashow, M.F. (1997). *Arabidopsis thaliana* CBF1 encodes an AP2 domain-containing transcriptional activator that binds to the C-repeat/DRE, a cis-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit. Proc. Natl. Acad. Sci. USA 94, 1035-1040.

Taylor, R.L. (1967). The foliar embryos of *Malaxis paludosa*. Can. J. Bot. 45, 1553-1556.

Taylor, N. G., Scheible, W. R., Cutler, S., Somerville, C. R., and Turner, S. R. (1999). The irregular xylem3 locus of *Arabidopsis* encodes a cellulose synthase required for secondary cell wall synthesis. Plant cell 11, 769-779.

Treacy, B.K., Hattori, J., Prud'homme, I., Barbour, E., Boutilier, K., Baszczynski, C.L., Huang, B., Johnson, D.A. and Miki, B.L. (1997). Bnm1, a Brassica pollen-specific gene. Plant Mol. Biol. 34, 603-611.

Twell, D., Wing, R., Yamaguchi, J. and McCormick, S. (1989). Isolation and expression of an anther-specific gene from tomato. Mol. Gen. Genet. 217, 240-245.

Ulmasov, T., Murfett, J., Hagen, G. and Guilfoyle, T. (1997). Aux/IAA proteins repress expression of reporter genes containing natural and highly active synthetic auxin response elements. Plant Cell 9, 1963-1971.

van Engelen, F.A., Molthoff, J.W., Conner, A.J., Nap, J.P., Pereira, A. and Stiekema, W.J. (1995). pBINPLUS: an improved plant transformation vector based on pBIN19. Transgenic Res. 4, 288-290.

Vergunst, A.C., de Waal, E.C. and Hooykaas, P.J.J. (1998). Root transformation by *Agrobacterium tumefaciens*. In *Arabidopsis Protocols*, J.M. Martinez-Zapater and J. Salinas, eds (Totowa, USA: Humana Press).

Xu, D., McElroy, D., Thornburg, R.W. and Wu, R.C.S. (1993). Systemic induction of a potato pin2 promoter by wounding, methyl jasmonate, and abscisic acid in transgenic rice plants. Plant Mol. Biol. 22, 573-588.

Xu, H., Knox, R.B., Taylor, P.E. and Singh, M.B. (1995). Bcp1, a gene required for male fertility in Arabidopsis. Proc. Natl. Acad. Sci. USA 92, 2106-2110.

Yarbrough, J.A. (1932). Anatomical and developmental studies of the foliar embryos of Bryophyllum calycinum. Amer. J. Bot. 19, 443-453.

Yeung, E.C. (1995). Structural and developmental patterns in somatic embryogenesis. In In Vitro Embryogenesis in Plants, T.A. Thorpe, ed (Dordrecht: Kluwer Academic Publishers), pp. 205-247.

Zarsky, V., Garrido, D., Eller, N., Tupy, J., Vicente, O., Schffl, F. and Heberle-Bors, E. (1995). The expression of a small heat shock gene is activated during induction of tobacco pollen embryogenesis by starvation. Plant Cell Environ. 18, 139-147.

Zhou, J., Tang, X. and Martin, G.B. (1997). The Pto kinase conferring resistance to tomato bacterial speck disease interacts with proteins that bind a cis-element of pathogenesis-related genes. EMBO J. 16, 3207-3218.

**THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE
PROPERTY OF PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:**

1. An isolated DNA molecule comprising a nucleotide sequence that hybridizes to the nucleotide sequence selected from the group consisting of SEQ ID NO's:1, 3 and 5 or a fragment or derivative thereof, excluding an AP2 domain repeat1-linker-AP2 domain repeat2 region, under moderate or stringent hybridization conditions.
2. The isolated DNA molecule of claim 1 wherein said isolated DNA molecule comprises at least 27 contiguous nucleotides of a nucleotide sequence selected from the group consisting of SEQ ID NO:1, 3 and 5.
3. The isolated DNA molecule of claim 1 wherein said isolated DNA molecule comprises a nucleotide sequence that is at least 70% homologous with a nucleotide sequence, or a fragment or derivative thereof, selected from the group consisting of SEQ ID NO:1, 3 and 5.
4. An isolated DNA molecule comprising a nucleic acid sequence encoding a protein, wherein said protein when present at a sufficient level within a plant cell renders said cell embryogenic, increases the regenerative capacity of said plant cell, or both renders said plant cell embryogenic and increases the regenerative capacity of said plant cell, said isolated DNA molecule having at least 70% homology within a nucleotide sequence, or a fragment or derivative thereof, selected from the group consisting of SEQ ID NO's:1, 3 and 5.
5. The isolated DNA molecule of claim 4 comprising a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5 or a fragment or derivative thereof, under moderate or stringent conditions.

6. The isolated DNA molecule of claim 4 comprising a nucleotide sequence that hybridizes to the nucleotide sequence of SEQ ID NO:1 or a fragment or derivative thereof, under moderate or stringent conditions.
7. The isolated DNA molecule of claim 4 comprising a nucleotide sequence that hybridizes to the nucleotide sequence of SEQ ID NO:3 or a fragment or derivative thereof, under stringent conditions.
8. The isolated DNA molecule of claim 6, wherein said DNA encodes a protein as defined by SEQ ID NO:2.
9. The isolated DNA molecule of claim 7, wherein said DNA encodes a protein as defined by SEQ ID NO:4.
10. A vector comprising the isolated DNA molecule as claimed in any one of claims 1 to 9, wherein said isolated DNA molecule is under control of a regulatory element that directs expression of said DNA in a plant cell.
11. The vector of claim 10, wherein said regulatory element is a constitutive regulatory element
12. The vector of claim 10, wherein said regulatory element is an inducible regulatory element.
13. The vector of claim 10, wherein said regulatory element is a tissue specific regulatory element
14. The vector of claim 10, wherein said regulatory element is an developmentally active regulatory element.
15. A transformed plant cell comprising the vector of any one of claims 10 to 14.

16. A transformed plant comprising the vector of any one of claims 10 to 14.
17. A seed obtained from the transformed plant of claim 16.
18. An isolated protein encoded by the isolated DNA molecule as claimed in any one of claims 4 to 9.
19. A method of producing asexually derived embryos comprising:
 - i) transforming a plant cell with the vector of any one of claims 10 to 14;
 - ii) growing said plant cell to produce transformed tissue;
 - iii) selecting said transformed tissue for occurrence of said isolated DNA molecule; and
 - iv) assaying said transformed plant for asexual embryo production.
20. The method of claim 19 wherein the step of assaying involves assaying for adventitious embryony.
21. The method of claim 19, wherein the step of assaying involves assaying for somatic embryos.
22. The method of claim 19, wherein the step of assaying involves assaying for gametophytic embryos.
23. The method of claim 19, wherein the step of assaying involves assaying for haploid parthenogenesis of the embryo sac.
24. The method of claim 19, wherein the step of assaying involves assaying for diplospory.

25. A method of modifying the regenerative capacity of a plant comprising
- i) transforming a plant cell with the vector of any one of claims 10 to 14;
 - ii) growing said transformed plant cell to produce transformed tissue; and
 - iii) assaying said transformed plant tissue for enhanced regeneration as compared to wild-type tissue.
26. The method of claim 25, wherein the step of growing said transformed plant cell, the step of assaying said transformed plant tissue, or both the step of growing said transformed plant cell and the step of assaying said transformed plant tissue are carried out in the absence of a growth regulator.
27. A method of selecting a transformed plant comprising;
- i) transforming a normally non-regenerative plant with a vector of any one of claims 10 to 14; and
 - ii) determining whether said transformed plant is able to regenerate under conditions in which said normally non-regenerative plant does not regenerate.
28. The isolated DNA molecule of claim 1 comprising a DNA sequence that comprises at least about 70% similarity with nucleotides 1-1619 of SEQ ID NO:5, or a fragment thereof.
29. The isolated DNA molecule of claim 1 wherein said isolated molecule comprises at least 22 contiguous nucleotides within nucleotides 1-1619 of SEQ ID NO:5.
30. A vector comprising the isolated DNA molecule of either claim 28 or 29 operably associated with a gene of interest, wherein said isolated DNA molecule directs the expression of said gene of interest within a plant cell.

31. The vector as defined by claim 30, wherein said gene of interest is heterologous with respect to the isolated DNA molecule.
32. The vector as defined by claim 31, wherein said gene of interest is selected from the group consisting of a pharmaceutically active protein, antibody, industrial enzyme, protein supplement, nutraceutical, storage protein, animal feed and animal feed supplement.
33. A transformed plant cell comprising the vector of either claim 30, 31 or 32.
34. A transformed plant comprising the vector of either claim 30, 31 or 32.
35. A seed obtained from the transformed plant of claim 34.
36. A method for directing the expression of a gene of interest within a developing embryo of a plant comprising transforming said plant with the vector as defined by either claim 30, 31 or 32.
37. A use of a nucleotide sequence as defined in any one of claims 4, 5, 6 or 7 as a selectable marker.
38. A method of producing asexually derived embryos comprising:
 - i) transiently transforming a plant cell with the vector of any one of claims 10 to 14, or introducing into said plant cell the protein of claim 18, to produce a modified plant cell;
 - ii) growing said modified plant cell to produce tissue; and
 - iii) assaying said tissue for asexual embryo formation.
39. The method of claim 38 wherein the step of assaying involves assaying for adventitious embryony.

40. The method of claim 38, wherein the step of assaying involves assaying for somatic embryos.
41. The method of claim 38, wherein the step of assaying involves assaying for gametophytic embryos.
42. The method of claim 38, wherein the step of assaying involves assaying for haploid parthenogenesis of the embryo sac.
43. The method of claim 38, wherein the step of assaying involves assaying for diplospory.
44. A method of modifying the regenerative capacity of a plant comprising
- i) transiently transforming a plant cell with the vector of any one of claims 10 to 14 or introducing into said plant cell the protein of claim 18, to produce a modified plant cell;
 - ii) growing said modified plant cell to produce tissue; and
 - iii) assaying said tissue for enhanced regeneration as compared to wild-type tissue.
45. The method of claim 44, wherein the step of growing said modified plant cell, the step of assaying said tissue, or both the step of growing said modified plant cell and the step of assaying said tissue are carried out in the absence of a growth regulator.
46. A method of producing an apomictic plant comprising:
- i) transforming a plant with the vector of any one of claims 10 to 14, to produce a transformed plant;
 - ii) selecting said transformed plant for occurrence of said isolated DNA molecule; and

- iii) assaying said transformed plant for asexual embryo production.
47. The method of claim 46 wherein the step of assaying involves assaying for adventitious embryony.
48. The method of claim 46, wherein the step of assaying involves assaying for somatic embryos.
49. The method of claim 46, wherein the step of assaying involves assaying for gametophytic embryos.
50. The method of claim 46, wherein the step of assaying involves assaying for parthenogenesis of the embryo sac.
51. A method of modifying the regenerative capacity of a plant comprising
- i) transiently transforming a plant cell with the vector of any one of claims 10 to 14, or introducing into said plant cell the protein of claim 18;
 - ii) growing said plant cell to form tissue; and
 - iii) assaying said tissue for enhanced regeneration as compared to wild-type tissue.
52. The method of claim 51, wherein the step of growing said plant cell, the step of assaying said tissue, or both the step of growing said plant cell and the step of assaying said tissue are carried out in the absence of a growth regulator.
53. A method of selecting a modified plant comprising;
- i) transiently transforming a normally non-regenerative plant with a vector of any one of claims 10 to 14, or introducing into said normally non-regenerative plant the protein of claim 18, to produce said modified plant; and

- ii) determining whether said modified plant is able to regenerate under conditions in which said normally non-regenerative plant does not germinate.
54. An isolated DNA molecule comprising a sequence encoding a protein consisting of two AP2 DNA binding domains, which when said protein is expressed at a sufficient level in a plant cell, renders said cell embryogenic, or increase the regenerative capacity of said plant cell, or both renders said cell embryogenic and increase the regenerative capacity of said plant cell.
55. A method of producing a protein of interest comprising
- i) transforming a plant with at least one vector, said at least one vector selected from any one of claims 10 to 14 to produce a transformed plant;
 - ii) selecting said transformed plant for occurrence of said isolated DNA molecule; and
 - iv) growing said transformed plant in order to produce said protein of interest, wherein expression of said protein of interest is induced by the expression product of said isolated DNA.
56. The method of claim 55, wherein said transformed plant is transformed with a second vector comprising a nucleotide sequence encoding said protein of interest under the control of a regulatory element, said regulatory element induced by the expression product of said isolated DNA..
57. The method of claim 55, wherein said protein of interest is a native protein.
58. The method of any one of claims 55 or 56, wherein said protein of interest is selected from the group consisting of a pharmaceutically active protein, antibody, industrial enzyme, protein supplement, nutraceutical, storage protein, an enzyme involved in oil biosynthesis, animal feed, and animal feed supplement.

59. The isolated DNA molecule of claim of any one of claims 4 to 7, wherein said isolated DNA molecule encodes a protein that is at least 70% similar with the amino acid defined by SEQ ID NO:2.
60. The isolated DNA molecule of claim of any one of claims 4 to 7, wherein said isolated DNA molecule encodes a protein that is at least 70% similar with the amino acid defined by SEQ ID NO:4.
61. The isolated protein of claim 18, wherein said protein comprises from about 30 to about 541 amino acids of the sequence disclosed in SEQ ID NO:2
62. The isolated protein of claim 18, wherein said protein comprises from about from about 30 to about 561 amino acids of the sequence disclosed in SEQ ID NO: 4.
63. An isolated DNA molecule comprising a nucleotide sequence that hybridizes to SEQ ID NO:6, excluding an AP2 domain repeat1-linker-AP2 domain repeat2 region, under stringent conditions.
64. The isolated DNA molecule of claim 4 comprising a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:6 or a fragment or derivative thereof, under moderate or stringent conditions.
65. The isolated DNA molecule of claim 64, wherein said DNA encodes a protein as defined by SEQ ID NO:7.
66. A vector comprising the isolated DNA molecule as claimed in any one of claims 63 to 65, wherein said isolated DNA molecule is under control of a regulatory element that directs expression of said DNA in a plant cell.
67. The vector of claim 66, wherein said regulatory element is a constitutive regulatory element

68. The vector of claim 67, wherein said regulatory element is an inducible regulatory element.
69. The vector of claim 67, wherein said regulatory element is a tissue specific regulatory element
70. The vector of claim 67, wherein said regulatory element is an developmentally active regulatory element.
71. A transformed plant cell comprising the vector of any one of claims 66 to 70.
72. A transformed plant comprising the vector of any one of claims 66 to 70.
73. A seed obtained from the transformed plant of claim 72.
74. An isolated protein encoded by the isolated DNA molecule as claimed in claim 65.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 December 2000 (14.12.2000)

PCT

(10) International Publication Number
WO 00/75330 A1

(51) International Patent Classification: C12N 15/29,
15/82, 5/10, C07K 14/415, A01H 5/00, 5/10

(21) International Application Number: PCT/CA00/00642

(22) International Filing Date: 2 June 2000 (02.06.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
99201745.9 2 June 1999 (02.06.1999) EP

(71) Applicants (for all designated States except US): **PLANT RESEARCH INTERNATIONAL (NL/NL)**; Building No. 107, Droeveendaalsesteeg 1, NL-6708 PB Wageningen (NL). **HER MAJESTY THE QUEEN IN RIGHT OF CANADA**, as represented by **THE MINISTER OF AGRICULTURE AND AGRI-FOOD CANADA [CA/CA]**; Eastern Cereal & Oilseed Research Centre, K.W. Neaby Building, Ottawa, Ontario K1A 0C6 (CA).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BOUTILIER**,

Kim [CA/NL]; Groen van Prinstererstraat 87, NL-6702 CP Wageningen (NL). **OUELLET, Thérèse [CA/CA]**; 115 Kennevale Drive, Nepean, Ontario K2J 3X7 (CA). **CUSTERS, Jan [NL/NL]**; Gen. Foulkesweg 25, NL-6703 BL Wageningen (NL). **HATTORI, Jiro [JP/CA]**; 763 Halstead Street, Ottawa, Ontario K1G 1M5 (CA). **MIKI, Brian [CA/CA]**; 1876 Dorset Drive, Ottawa, Ontario K1H 5V1 (CA). **VAN LOOKEREN CAMPAGNE, Michiel [NL/BE]**; Lievestraat 31, B-9920 Lovendegem (BE).

(74) Agents: **SECHLEY, Konrad, A. et al.**; Gowling Lafleur Henderson LLP, Suite 2600, 160 Elgin Street, Ottawa, Ontario K1P 1C3 (CA).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian

[Continued on next page]

(54) Title: **USE OF THE BNM3 TRANSCRIPTIONAL ACTIVATOR TO CONTROL PLANT EMBRYOGENESIS AND REGENERATION PROCESSES**

In vivo development

In vitro development



(57) Abstract: The present invention provides for a gene obtained during the induction of microspore embryogenesis. The protein encoded by this gene renders plant cells embryogenic, and increases the regenerative capacity of the plant cell. Also disclosed is the regulatory region of this gene and its use for directing the expression of a gene of interest within a suitable host cell.




FIGURE 1

FIGURE 2

BNM3A GTTCATCTCTCTTCTTTAAGACCAAACCTTTTCTCCTCCTCTTCATGCATGAACCTTA 60
BNM3B -----

BNM3A ACTAAGTTCCTTCTCTTTTACCTTTTACCAAGAAGCTCGTTAGATCACTCTCTGAAGCTCAA 120
BNM3B -----TCTTCTTTTACCTTTTACCAAGAAGCTCGTTAGATCATTTTCTGAAGCTCGA 51

BNM3A TGAATAATAACTGGTTAGGCTTTTCTCTCTCTCCTTATGAACAAAATCACCATCGTAAGG 180
BNM3B TGAATAATAACTGGTTAGGCTTTTCTCTCTCTCCTTATGAACAAAATCACCATCGTAAGG 111

BNM3A ACGTCTACTCTTCACCACCAACCGTCGTAGATGTCGCGGAGAGTACTGTTACGATC 240
BNM3B ACGTCTCTCTTCCACCACCAACCGCGGTAGATGTCGCGGAGAGTACTGTTACGATC 171

BNM3A CGACCGCTGCCTCCGATGAGTCTTCAGCCATCCAAACATCGTTTCTCTCTCCCTTTGGTG 300
BNM3B CGACCGCTGCCTCCGATGAGTCTTCAGCCATCCAAACATCGTTTCTCTCTCCCTTTGGTG 231

BNM3A TCGTCGTTCGATGCTTTTCCACGAGACAAACAATAGTCACTCCGAGATTGGGACATCAATG 360
BNM3B TCGTCCTCGATGCTTTTCCACGAGACAAACAATAGTCACTCCGAGATTGGGACATCAATG 291

BNM3A GTTGTGCATGCAATAACATCCACAACGATGAGCAAGATGGACCAAAGCTTGAGAAATTTCC 420
BNM3B GTAGTGCATGTAATAACATCCACAATGATGAGCAAGATGGACCAAACCTTGAGAAATTTCC 351

BNM3A TTGGCCGACACCAACGATTTTCAACACCAACGAAACGTTGGAGATGGAAGTGGAAAGTG 480
BNM3B TTGGCCGACACCAACGATTTTCAACACCAACGAAACGTTGGAGATATCGATGGAAGTG 411

BNM3A GCTGTTATGGAGAGGAGACGGTGGTGGTGGCTCACTAGGACTTTGATGATAAAGACAT 540
BNM3B GGTGTTATGGAGAGGAGACGGTGGTGGTGGCTCACTAGGACTTTGATGATAAAGACAT 471

BNM3A GGCTGAGAAATCAACCCGTGGATAATGTTGATAATCAAGAAATGGCAATGCTGCAAAAG 600
BNM3B GGCTGAGAAATCAACCCGTGGATAATGTTGATAATCAAGAAATGGCAATGCTGCAAAAG 531

BNM3A GCCTGTCCCTCTCAATGAAGTCTCATCTACTCTTGTGTATACCAACACGACAGCAATAACA 660
BNM3B GCCTGTCCCTCTCAATGAAGTCTCATCTACTCTTGTGTATACCAACACGACAGCAATAACA 591

BNM3A ACGTTGTTGCCCAAGGGAAGACTATTGATGATAGCGTTGAAGCTACACCGAAGAAAACCTA 720
BNM3B ACCCTGTTGCCCAAGGGAAGACTATTGATGATAGCGTTGAAGCTACACCGAAGAAAACCTA 651

BNM3A TTGAGAGTTTGGACAGAGGACGTCATATACCGCGGTGTTACRAGGCATCGGTGGACAG 780
BNM3B TTGAGAGTTTGGACAGAGGACGTCATATACCGCGGTGTTACRAGGCATCGGTGGACAG 711

BNM3A GAAGATATGAGGCACATTTATGGGATAATAGTTGTAAAGAGAAGGCCAAACGCGCAAG 840
BNM3B GAAGATATGAGGCACATTTATGGGATAATAGTTGTAAAGAGAAGGCCAAACGCGCAAG 771

BNM3A GAAGACAAGTTTATTGGGAGGTTATGACAAAGAAGAAAAGCAGCTAGGGCTTATGATT 900
BNM3B GAAGACAAGTTTATTGGGAGGTTATGACAAAGAAGAAAAGCAGCTAGGGCTTATGATT 831

20000364-140000

FIGURE 2(Cont'd)

BNM3A	TAGCCGCACTCAAGTATTGGGGAACCACTACTACTAACTTCCCATGAGCGAATATG	960
BNM3B	TAGCCGCACTCAAGTATTGGGGAACCACTACTACTAACTTCCCATGAGCGAATATG	891
BNM3A	AAAAAGAGGTAGAAGAGATGAAGCACATGACAAGGCAAGAGTATGTTGCCCTCACTGCGCA	1020
BNM3B	AGAAAGAGATAGAAGAGATGAAGCACATGACAAGGCAAGAGTATGTTGCCCTCACTGCGCA	951
BNM3A	GGAAAAAGTAGTGGTTTCTCTCGTGGTGTCATCGATTATCTGTGGAGTAACAAGACATCACC	1080
BNM3B	GGAAAAAGTAGTGGTTTCTCTCGTGGTGTCATCGATTATCTGTGGAGTAACAAGACATCACC	1011
BNM3A	AACATGGAAGATGGCAAGCTAGGATAGGAAGAGTGCCTGGTAACAAGACCTCTACTTTGG	1140
BNM3B	AACATGGAAGATGGCAAGCTAGGATAGGAAGAGTGCCTGGTAACAAGACCTCTACTTTGG	1071
BNM3A	GAACCTTTGGGCACACAGAAGAAGCTGCAGAGGCATACGACATTGCGGCCATCAAATTCA	1200
BNM3B	GAACCTTTGGGCACACAGAAGAAGCTGCAGAGGCATACGACATTGCGGCCATCAAATTCA	1131
BNM3A	GAGGATTAACCGCAGTGACTAACTTCGACATGAACAGATACAAAGCTTAAAGCAATCCTCG	1260
BNM3B	GAGGATTAACCGCAGTGACTAACTTCGACATGAACAGATACAAAGCTTAAAGCAATCCTCG	1191
BNM3A	AAAGCCCTAGTCTTCTTATTGGTAGCGCGCAAAACGCTCTCAAGGAGGCTAACCGTCCGG	1320
BNM3B	AAAGCCCTAGTCTTCTTATTGGTAGCGCGCAAAACGCTCTCAAGGAGGCTAACCGTCCGG	1251
BNM3A	TTCCAAAGTATGATGATGATCAGTAATAACGTTTCAGAGAGTGAAGATAGTCTAGCGGTT	1380
BNM3B	TTCCAAAGTATGATGATGATCAGTAATAACGTTTCAGAGAGTGAAGATAGTCTAGCGGTT	1311
BNM3A	GGCAAAACGCTGCGGTTTCAGCATCATCAGGAGTAGATTGAGCTTATTGCACCAACATC	1440
BNM3B	GGCAAAACGCTGCGGTTTCAGCATCATCAGGAGTAGATTGAGCTTATTGCACCAACATC	1371
BNM3A	AAGAGAGGTACAATGGTTATTATACAATGGAGGAACTTGTCTTCGGAGAGTGCTAGGG	1500
BNM3B	AAGAGAGGTACAATGGTTATTATACAATGGAGGAACTTGTCTTCGGAGAGTGCTAGGG	1431
BNM3A	CTTGTTTCAAAACAGAGGATGATCAACACCATTTCTTGAGCAACACGAGAGCCTCATGA	1560
BNM3B	CTTGTTTCAAAACAGAGGATGATCAACACCATTTCTTGAGCAACACGAGAGCCTCATGA	1491
BNM3A	CTAATATCGATCATCAAAGTTCTGTTTCGAGATGATTCGGTTACTGTTTGTGGAATGTTG	1620
BNM3B	CTAATATCGATCATCAAAGTTCTGTTTCGAGATGATTCGGTTACTGTTTGTGGAATGTTG	1551
BNM3A	TTGGTTATGGTGGTTATCAAGGATTTCAGCCCCGGTTAACTGCGATGCCCTACGCTGCTA	1680
BNM3B	TTGGTTATGGTGGTTATCAAGGATTTCAGCCCCGGTTAACTGCGATGCCCTACGCTGCTA	1611
BNM3A	GTGAGTTTGATTATACCGCAAGAAACCAATTATTACTTTGCTCAGCAGCAGCAGCCGAGC	1740
BNM3B	GTGAGTTTGATTATACCGCAAGAAACCAATTATTACTTTGCTCAGCAGCAGCAGCCGAGC	1671
BNM3A	AGTCGCCAGGTGGAGATTTTCCCGCGGCAATGACGAATAATGTTGGCTCTAATATGTATT	1800
BNM3B	ATTCGCCAGGAGGAGATTTTCCCGCGGCAATGACGAATAATGTTGGCTCTAATATGTATT	1731
BNM3A	ACCATGGGGAAGGTGGTGGAGAAGTTGCTCCAACATTTACAGTTTGGAAACGACAAATTAGA	1860
BNM3B	ACCATGGGGAAGGTGGTGGAGAAGTTGCTCCAACATTTACAGTTTGGAAACGACAAATTAGA	1791

000001-000001

FIGURE 2(Cont'd)

BNM3A	AAAAATAGTTAAAGATCTTTAGTTATATGCGTTGTTGTGTGCTGGTGAACAGTGTGATAC	1920
BNM3B	AATAATAGTTAAAGATCTTTAGTTATATGCGTTGTTGTGTGCTGGTGAACAGTTTGATAC	1851
	** *****	
BNM3A	TTTGATTATGTTTTTCTTTCTTTTCCTTTTCTTGGTTAATTTCTTAAGACTTATT	1980
BNM3B	TTTGATTATGTTTTT--TTTCTCTTTTCATTTTGTGGTTAGTTTCTTAAGACTTATT	1909

BNM3A	TTTAGTTTCCATTAGTTGGATAAAATTTTCAGACT-----	2014
BNM3B	TTTGTTTTCCATTAGTTGGATAAAATTTTCGGACTTAAGGGTCACCTTCTGTTCTGACTTCT	1969

BNM3A	-----	
BNM3B	GTCTAATACAGAAAAGTTTTTCAT	1992

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095

5/11

FIGURE 3

```

BNM3A MNNWLGFSLSPYEQNHHRKDVYSSTTTTVVVDVAGEYCYDPTAASDESSAIQTSFFSPFG 60
BNM3B MNNWLGFSLSPYEQNHHRKDVCSSTTTTAVDVAGEYCYDPTAASDESSAIQTSFFSPFG
*****

BNM3A VVVDFAFTRDNNSHSRDWDINGCACCNNIHNDQDQGPKEFLGRITTTIYNTNENVGDGSGS 120
BNM3B VVLDFAFTRDNNSHSRDWDINGSACCNNIHNDQDQGPKEFLGRITTTIYNTNENVGDIDGS
*****

BNM3A GCGGGDGGGSLGLSMIKTWLRNOPVDNVNDQENGNAAGLSLSMNSSTSDDNNNSNN 180
BNM3B GCGGGDGGGSLGLSMIKTWLRNOPVDNVNDQENGNAAGLSLSMNSSTSDDNNNYSSN
*****

                                repeat 1
BNM3A NVVAQGKTIDDSVEATPKKTIIESFGQRTSIYRGVTRHRWTGRYEAHLWDNSCKREGQTRK 240
BNM3B NLVAQGKTIDDSVEATPKKTIIESFGQRTSIYRGVTRHRWTGRYEAHLWDNSCKREGQTRK
*****

BNM3A GRQVYLGGYDKKEKAARAYDLAALKYWGTTTTTNFFMSEYEKEVEEMKHMTRQEYVASLR 300
BNM3B GRQVYLGGYDKKEKAARAYDLAALKYWGTTTTTNFFMSEYEKEIEEMKHMTRQEYVASLR
*****

                                repeat 2
BNM3A RKSSGFSRGASIYRGVTRHEQHGRWQARIGRVAGNKDLYLGTGTQEAAEAYDLAAIKF 360
BNM3B RKSSGFSRGASIYRGVTRHEQHGRWQARIGRVAGNKDLYLGTGTQEAAEAYDLAAIKF
*****

BNM3A RGLTAVTNFDMNRYNVKAILSPSLPIGSAAKRLKEANRPVPSMMMISSNNVSESENSASG 420
BNM3B RGLTAVTNFDMNRYNVKAILSPSLPIGSAAKRLKEANRPVPSMMMISSNNVSEENNASG
*****


BNM3A WQNAAVQHHCQGVDSLHLHQHQRERYNGYYNGGNLSSSARACFKQEDDQHHFLSNTQSLM 480
BNM3B WQNAAVQHHCQGVDSLHLHQHQRERYNGYYNGGNLSSSARACFKQEDDQHHFLSNTQSLM
*****

BNM3A TNIDHQSSVSDSVTCGNVVGYYGGYQGFAPVNCDAAYAASEFDYNARNHYFAQQQQQTQ 540
BNM3B TNIDHQSSVSDSVTCGNVVGYYGGYQGFAPVNCDAAYAASEFDYNARNHYFAQQQQQTQ
*****

BNM3A QSPGGDFPAAMTNNVGSNNMYHGGEGGEVAPTFTVWVNDN 579
BNM3B HSPGGDFPAAMTNNVGSNNMYHGGEGGEVAPTFTVWVNDN
*****

```

Figure 4



208040.42908660

WO 00/75330

Inventor: BOUTILIER et al.
 Docket No.: 270.6213/VO
 Title: USE OF THE BNM3 TRANSCRIPTIONAL ACTIVATOR TO CONTROL PLANT
 EMBRYO GENESIS AND REGENERATION PROCESS
 Attorney Name: Gregory A. Sebald
 Phone No.: 612.336 4728
 Sheet 7 of 11

PCT/CA00/00642

09/980364

FIGURE 5

AP2 DOMAIN REPEAT 1

BNM3 TSYRGVTRHRWTCRVEAHJWNSCKREGQTRKGRQVYLGGYDKEKAARAYDLAALKYWGTTTTTTFPMSEYEKEV
 ANT T*Q*****F*K*HS*****M*****PS*H**SHEN*Q**I
 ZM T*****E**R*****S*****F**P*****QV*N*****L
 GL15 S*Q*****FY*R**W*S*I*--C-----*K*****F*TAQA*****Q*I*FR*LNADI**TLDD*KD*M
 AP2 S*Q*****FY*R**W*S*I*--C-----*K*****F*TAHA*****R*I*FR*VEADI**NIDD*DDDL


LINKER

BNM3 EEMKHMTRQEVVASLRKSSGFSRG
 ANT *D*N*****H*****
 ZM ***S*****F*****V**
 GL15 KK**DLSKE*P*LV**QGR**V**
 AP2 KQ*TNL*KE*F*HV**Q*I**P**

AP1 DOMAIN REPEAT 2

BNM3 ASYRGVTRHHQHGRWQARIGRVAGNKDLYLGTGTQEEAAEAYDIAAIKFRGLTAVTNFDMNRYNVKA
 ANT *****S*****V*****TN*****IT*D*DR
 ZM *****S*****N*****L*S*D*ES
 GL15 S*RP***-Q*KC*K*E***QLM*K*YV**LYD*ET***Q***K***CY*KE*****AQ*S*DKEL
 AP2 S*K*****L*KC***E**M*QFL*K*YV**L*D*EV***R**K***CN*KD*****PSI*DEEL


FIGURE 6



WO 00/75330


CA00/00642

FIGURE 7



10/11

FIGURE 8




09/980364

WO 00/75330


PCT/CA00/00642

FIGURE 9

A



B





MERCHANT & GOULD P.C.

United States Patent Application

COMBINED DECLARATION AND POWER OF ATTORNEY

I, , named inventor I hereby declare that: my residence, post office address and citizenship are as stated below next to my name; that

I verily believe I am the original, first and sole inventor (if only one name is listed below) or a joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: **USE OF THE BNM3 TRANSCRIPTIONAL ACTIVATOR TO CONTROL PLANT EMBRYOGENESIS AND REGENERATION PROCESS**

The specification of which

- a. ☐ is attached hereto
 b. ☒ was filed on as application serial no. and was amended on (if applicable) (in the case of a PCT-filed application) described and claimed in international no. PCT/CA00/00642 filed June 2, 2000 and as amended on (if any), which I have reviewed and for which I solicit a United States patent.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I hereby claim foreign priority benefits under Title 35, United States Code, § 119/365 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on the basis of which priority is claimed:

- a. ☐ No such applications have been filed.
 b. ☒ Such applications have been filed as follows:

FOREIGN APPLICATION(S), IF ANY, CLAIMING PRIORITY UNDER 35 USC § 119			
COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	DATE OF ISSUE (day, month, year)
ESP	99201745.9	2 June 1999	
ALL FOREIGN APPLICATION(S), IF ANY, FILED BEFORE THE PRIORITY APPLICATION(S)			
COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	DATE OF ISSUE (day, month, year)

I hereby claim the benefit under Title 35, United States Code, § 120/365 of any United States and PCT international application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, § 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. APPLICATION NUMBER	DATE OF FILING (day, month, year)	STATUS (patented, pending, abandoned)

I hereby claim the benefit under Title 35, United States Code § 119(e) of any United States provisional application(s) listed below:

U.S. PROVISIONAL APPLICATION NUMBER	DATE OF FILING (Day, Month, Year)

I acknowledge the duty to disclose information that is material to the patentability of this application in accordance with Title 37, Code of Federal Regulations, § 1.56 (reprinted below):

§ 1.56 Duty to disclose information material to patentability.

(a) A patent by its very nature is affected with a public interest. The public interest is best served, and the most effective patent examination occurs when, at the time an application is being examined, the Office is aware of and evaluates the teachings of all information material to patentability. Each individual associated with the filing and prosecution of a patent application has a duty of candor and good faith in dealing with the Office, which includes a duty to disclose to the Office all information known to that individual to be material to patentability as defined in this section. The duty to disclose information exists with respect to each pending claim until the claim is canceled or withdrawn from consideration, or the application becomes abandoned. Information material to the patentability of a claim that is canceled or withdrawn from consideration need not be submitted if the information is not material to the patentability of any claim remaining under consideration in the application. There is no duty to submit information which is not material to the patentability of any existing claim. The duty to disclose all information known to be material to patentability is deemed to be satisfied if all information known to be material to patentability of any claim issued in a patent was cited by the Office or submitted to the Office in the manner prescribed by §§ 1.97(b)-(d) and 1.98. However, no patent will be granted on an application in connection with which fraud on the Office was practiced or attempted or the duty of disclosure was violated through bad faith or intentional misconduct. The Office encourages applicants to carefully examine:

(1) prior art cited in search reports of a foreign patent office in a counterpart application, and

(2) the closest information over which individuals associated with the filing or prosecution of a patent application believe any pending claim patentably defines, to make sure that any material information contained therein is disclosed to the Office.

(b) Under this section, information is material to patentability when it is not cumulative to information already of record or being made of record in the application, and

(1) It establishes, by itself or in combination with other information, a prima facie case of unpatentability of a claim;

(2) It refutes, or is inconsistent with, a position the applicant takes in:

(i) Opposing an argument of unpatentability relied on by the Office, or

(ii) Asserting an argument of patentability.

A prima facie case of unpatentability is established when the information compels a conclusion that a claim is unpatentable under the preponderance of evidence, burden-of-proof standard, giving each term in the claim its broadest reasonable construction consistent with the specification, and before any consideration is given to evidence which may be submitted in an attempt to establish a contrary conclusion of patentability.

(c) Individuals associated with the filing or prosecution of a patent application within the meaning of this section are:

(1) Each inventor named in the application;

(2) Each attorney or agent who prepares or prosecutes the application; and

(3) Every other person who is substantively involved in the preparation or prosecution of the application and who is associated with the inventor, with the assignee or with anyone to whom there is an obligation to assign the application.

(d) Individuals other than the attorney, agent or inventor may comply with this section by disclosing information to the attorney, agent, or inventor.

(e) In any continuation-in-part application, the duty under this section includes the duty to disclose to the Office all information known to the person to be material to patentability, as defined in paragraph (b) of this section, which became available between the filing date of the prior application and the national or PCT international filing date of the continuation-in-part application.

I hereby appoint the following attorney(s) and/or patent agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected herewith:

Albrecht, John W. Reg. No. 40,481
Ali, M. Jeffer Reg. No. 46,359
Alter, Allan G. Reg. No. 40,274
Anderson, Gregg I. Reg. No. 28,828
Batzli, Brian H. Reg. No. 32,960
Beard, John L. Reg. No. 27,612
Berns, John M. Reg. No. 43,496
Branch, John W. Reg. No. 41,633
Brown, Jeffrey C. Reg. No. 41,643
Bruess, Steven C. Reg. No. 34,130
Byrne, Linda M. Reg. No. 32,404
Campbell, Keith Reg. No. 46,597
Carlson, Alan G. Reg. No. 25,959
Caspers, Philip P. Reg. No. 33,227
Clifford, John A. Reg. No. 30,247
Cook, Jeffrey Reg. No. 48,649
Daigault, Ronald A. Reg. No. 25,968
Daley, Dennis R. Reg. No. 34,994
Daulton, Julie R. Reg. No. 36,414
DeVries Smith, Katherine M. Reg. No. 42,157
DiPietro, Mark J. Reg. No. 28,707
Dose, Matthew A. Reg. No. P-48,957
Edell, Robert T. Reg. No. 20,187
Epp, Ryan, Sandra Reg. No. 39,667
Glance, Robert J. Reg. No. 40,620
Goff, Jared S. Reg. No. 44,716
Goggin, Matthew J. Reg. No. 44,125
Golla, Charles E. Reg. No. 26,896
Gorman, Alan G. Reg. No. 38,472
Gould, John D. Reg. No. 18,223
Gregson, Richard Reg. No. 41,804
Gresens, John J. Reg. No. 33,112
Hammer, Samuel A. Reg. No. 46,754
Hammer, Curtis B. Reg. No. 29,165
Harrison, Kevin C. Reg. No. 46,759
Hertzberg, Brett A. Reg. No. 42,660
Hillson, Randall A. Reg. No. 31,838
Holzer, Jr., Richard J. Reg. No. 42,668
Hope, Leonard J. Reg. No. 44,774
Jardine, John S. Reg. No. P-48,835
Johns, Nicholas P. Reg. No. 48,995
Johnston, Scott W. Reg. No. 39,721
Kadievitch, Natalie D. Reg. No. 34,196
Kaseburg, Frederick A. Reg. No. 47,695
Kettelberger, Denise Reg. No. 33,924
Keys, Jeramie J. Reg. No. 42,724
Knearl, Homer L. Reg. No. 21,197
Kowalchuk, Alan W. Reg. No. 31,535
Kowalchuk, Katherine M. Reg. No. 36,848
Lacy, Paul E. Reg. No. 38,946
Larson, James A. Reg. No. 40,443

100

Leonard, Christopher J. Reg. No. 41,940
Liepa, Mara E. Reg. No. 40,066
Lindquist, Timothy A. Reg. No. 40,701
Lown, Jean A. Reg. No. 48,428
Mayfield, Denise L. Reg. No. 33,732
McDonald, Daniel W. Reg. No. 32,044
McIntyre, Jr., William F. Reg. No. 44,921
Mitchem, M. Todd Reg. No. 40,731
Mueller, Douglas P. Reg. No. 30,300
Nelson, Anna M. Reg. No. 48,935
Paley, Kenneth B. Reg. No. 38,989
Parsons, Nancy J. Reg. No. 40,364
Pauly, Daniel M. Reg. No. 40,123
Phillips, John B. Reg. No. 37,206
Pino, Mark J. Reg. No. 43,858
Prendergast, Paul Reg. No. 46,068
Pytel, Melissa J. Reg. No. 41,512
Qualey, Terry Reg. No. 25,148
Reich, John C. Reg. No. 37,703
Reiland, Earl D. Reg. No. 25,767
Samuels, Lisa A. Reg. No. 43,080
Schmaltz, David G. Reg. No. 39,828
Schuman, Mark D. Reg. No. 31,197
Schumann, Michael D. Reg. No. 30,422
Scull, Timothy B. Reg. No. 42,137
Sebald, Gregory A. Reg. No. 33,280
Skoog, Mark T. Reg. No. 40,178
Spellman, Steven J. Reg. No. 45,124
Stewart, Alan R. Reg. No. 47,974
Stoll-DeBell, Kirstin L. Reg. No. 43,164
Sullivan, Timothy Reg. No. 47,981
Summer, John P. Reg. No. 29,114
Swenson, Erik G. Reg. No. 45,147
Tellekson, David K. Reg. No. 32,314
Trembath, Jon R. Reg. No. 38,344
Tunheim, Marcia A. Reg. No. 42,189
Underhill, Albert L. Reg. No. 27,403
Vandenburgh, J. Derek Reg. No. 32,179
Wahl, John R. Reg. No. 33,044
Weaver, Paul L. Reg. No. 48,640
Welter, Paul A. Reg. No. 20,890
Whipps, Brian Reg. No. 43,261
Whitaker, John E. Reg. No. 42,222
Wier, David D. Reg. No. P-48,229
Williams, Douglas J. Reg. No. 27,054
Withers, James D. Reg. No. 40,376
Witt, Jonelle Reg. No. 41,980
Wong, Thomas S. Reg. No. 48,577
Wu, Tong Reg. No. 43,361
Young, Thomas Reg. No. 25,796
Zeuli, Anthony R. Reg. No. 45,255

I hereby authorize them to act and rely on instructions from and communicate directly with the person/assignee/attorney/firm/ organization who/which first sends/sent this case to them and by whom/which I hereby declare that I have consented after full disclosure to be represented unless/until I instruct Merchant & Gould P.C. to the contrary.

I understand that the execution of this document, and the grant of a power of attorney, does not in itself establish an attorney-client relationship between the undersigned and the law firm Merchant & Gould P.C., or any of its attorneys.

Please direct all correspondence in this case to Merchant & Gould P.C. at the address indicated below:

Merchant & Gould P.C.
P.O. Box 2903
Minneapolis, MN 55402-0903



I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

2	Full Name Of Inventor <u>1-00</u>	Family Name <u>Boutillier</u>	First Given Name <u>Kim</u>	Second Given Name
0	Residence & Citizenship	City <u>Wageningen</u>	State or Foreign Country <u>Netherlands</u>	Country of Citizenship <u>Canada</u> <u>CAX</u>
1	Mailing Address	Address <u>Green van Prinstererstraat 87</u>	City <u>Wageningen</u>	State & Zip Code/Country <u>NL-6702 CP / Netherlands</u>
Signature of Inventor 201: <u>Kim Boutillier</u>			Date: <u>11-01-02</u>	
2	Full Name Of Inventor <u>2-00</u>	Family Name <u>Ouellet</u>	First Given Name <u>Therese</u>	Second Given Name
0	Residence & Citizenship	City <u>Nepean</u>	State or Foreign Country <u>Ontario</u>	Country of Citizenship <u>Canada</u> <u>CAX</u>
2	Mailing Address	Address <u>115 Kennevale Drive</u>	City <u>Nepean</u>	State & Zip Code/Country <u>Ontario K2J 3X7 / Canada</u>
Signature of Inventor 202: <u>Therese Ouellet</u>			Date: <u>24-01-02</u>	
2	Full Name Of Inventor <u>3-00</u>	Family Name <u>Custers</u>	First Given Name <u>Jan</u>	Second Given Name
0	Residence & Citizenship	City <u>Wageningen</u>	State or Foreign Country <u>Netherlands</u>	Country of Citizenship <u>Netherlands</u> <u>NLX</u>
3	Mailing Address	Address <u>Gen Foulkestraat 25</u>	City <u>Wageningen</u>	State & Zip Code/Country <u>NL-6703 BL / Netherlands</u>
Signature of Inventor 203: <u>[Signature]</u>			Date:	
2	Full Name Of Inventor <u>4-00</u>	Family Name <u>Hattori</u>	First Given Name <u>Jiro</u>	Second Given Name
0	Residence & Citizenship	City <u>Ottawa</u>	State or Foreign Country <u>Ontario</u>	Country of Citizenship <u>Japan</u> <u>JPX</u>
4	Mailing Address	Address <u>753 Halstead Street</u>	City <u>Ottawa</u>	State & Zip Code/Country <u>Ontario K1G 1M5 / Canada</u>
Signature of Inventor 204: <u>Jiro Hattori</u>			Date: <u>2002-Jan 24</u>	
2	Full Name Of Inventor <u>5-00</u>	Family Name <u>Miki</u>	First Given Name <u>Brian</u>	Second Given Name
0	Residence & Citizenship	City <u>Ottawa</u>	State or Foreign Country <u>Ontario</u>	Country of Citizenship <u>Canada</u> <u>CAX</u>
5	Mailing Address	Address <u>1876 Dorset Drive</u>	City <u>Ottawa</u>	State & Zip Code/Country <u>Ontario K1H 5V1 / Canada</u>
Signature of Inventor 205: <u>[Signature]</u>			Date: <u>January 24, 2002</u>	



SEQUENCE LISTING

<110> Boutiller, Kim
Ouellet, Therese
Custers, Jan
Hattori, Jiro
Miki, Brian
Van Lookeren Campagne, Michiel

<120> Use of the BNM3 Transcriptional Activator to Control
Plant Embryogenesis and Regeneration Processes

<130> 270.62USWO

<140> 09/980,364

<141> Herewith

<150> EP 99201745.9-2106

<151> 1999-06-02

<160> 14

<170> PatentIn Ver. 2.1

<210> 1

<211> 2014

<212> DNA

<213> Brassica napus

<400> 1

gttcattctct cttctttaaag accaaaaacct ttttctctc ctcttcctgc atgaacccta 60
actaagttct tcttctttta ccttttacca agaactcgtt agatcactct ctgaactcaa 120
tgaataataa ctggttaggc ttttctctct ctcccttatga acaaaatcac catogtaagg 180
acgtctactc ttccaccacc acaaccgtcg tagatgtcgc cggagagtagc tgttaacgac 240
cgaccgctgc ctccgatgag tcttcagcca tccaaacatc gtttctctt ccttttggtg 300
tcgtcgtcga tgctttcacc agagacaaca atagtcactc ccgagattgg gacatcaatg 360
gtttgtcagt caataacatc cacaacgatg agcaagatgg accaaagctt gagaatttcc 420
ttggcgcaac caccacgatt tacaacacca acgaaaacgt tggagatgga agtggaagtg 480
gctgttatgg aggaggagac ggtggtggtg gctcactagg acttctgatg ataaagacat 540
ggctgagaaa tcaaccgctg gataatgttg ataatacaaga aaatggcaat gctgcaaaag 600
gcctgtccct ctcaatgaac tcactcactt cttgtgataa caacaacgac agcaataaca 660
acgttgttgc ccaaggggaag actattgatg atagcgttga agctacacgc aagaaaaacta 720
ttgagagtgt tggacagagg acgtctatat accgcggtgt tacaaggcat cgttgagacg 780
gaagatatga ggcacattta tgggataata gttgtaaaag agaaggccaa acgcgcgaag 840
gaagacaagt ttatttggga ggttatgaca aagaagaaaa agcagctagg gcttatgatt 900
tagccgcact caagtattgg ggaaccacca ctactactaa ctcccccatg agcgaatatg 960
aaaaagaggt agaagagatg aagcacatga caagggaaga gtatgttgc tcaactgcga 1020
ggaaaagtag tggtttctct cgtggtgcat cgatttatcg tggagttaaca agacataacc 1080
aacatggaag atggcaagct aggataggaa gagtgcgccg taacaaagac ctctacttgg 1140
gaacttttgg cacacaagaa gaagctgcag aggcatacga cattgcgcc atcaaatcca 1200
gaggattaac cgcagtgact aacttcgaca tgaacagata caacgttaaa gcaactctcg 1260
aaagccctag tcttctatt ggtagcgccg caaaacgtct caagaggagt aaccgctccg 1320
ttccaagtat gatgatgatc agtaataacg tttcagagag tgagaatagt gctagcggtt 1380
ggcaaaaacg tcgctgttcag catcatcagg gagttagatt gagcttattg caoccaacac 1440
aagagagtag caatggttat tattacaatg agggaactt gcttctcagg agtgctaggg 1500
cttgtttcaa acaagaggat gatcaacacc atttcttgag caacacgcag agcctcatga 1560

ctaataatcga tcatcaaagt tctgtttcgg atgattcgggt tactgttttgt ggaaatgttg 1620
 ttgggttatgg tgggttatcaa ggatttgcag ccccggttaa ctgcgatgcc tacgtctgcta 1680
 gtgagtttga ttataacgca agaaaccatt attactttgc tcagcagcag cagaccaccag 1740
 agtcgccagg tggagatttt ccgcgggcaa tgacgaataa tgttggtctc aatatgtatt 1800
 accatgggga agtggttgga gaagttgctc caacatttac agtttggaac gacaattaga 1860
 aaaaatagtt aaagatcttt agttatatgc gttgttgtgt gctgggtgaac agtgtgatac 1920
 tttgattatg tttttttctt tctctttttc tttttcttgg ttaatttctt aagacttatt 1980
 tttagtttcc attagttgga taaattttca gact 2014

<210> 2
 <211> 579
 <212> PRT
 <213> Brassica napus

<400> 2

Met Asn Asn Asn Trp Leu Gly Phe Ser Leu Ser Pro Tyr Glu Gln Asn
 1 5 10 15

His His Arg Lys Asp Val Tyr Ser Ser Thr Thr Thr Thr Val Val Asp
 20 25 30

Val Ala Gly Glu Tyr Cys Tyr Asp Pro Thr Ala Ala Ser Asp Glu Ser
 35 40 45

Ser Ala Ile Gln Thr Ser Phe Pro Ser Pro Phe Gly Val Val Val Asp
 50 55 60

Ala Phe Thr Arg Asp Asn Asn Ser His Ser Arg Asp Trp Asp Ile Asn
 65 70 75 80

Gly Cys Ala Cys Asn Asn Ile His Asn Asp Glu Gln Asp Gly Pro Lys
 85 90 95

Leu Glu Asn Phe Leu Gly Arg Thr Thr Thr Ile Tyr Asn Thr Asn Glu
 100 105 110

Asn Val Gly Asp Gly Ser Gly Ser Gly Cys Tyr Gly Gly Gly Asp Gly
 115 120 125

Gly Gly Gly Ser Leu Gly Leu Ser Met Ile Lys Thr Trp Leu Arg Asn
 130 135 140

Gln Pro Val Asp Asn Val Asp Asn Gln Glu Asn Gly Asn Ala Ala Lys
 145 150 155 160

Gly Leu Ser Leu Ser Met Asn Ser Ser Thr Ser Cys Asp Asn Asn Asn
 165 170 175

Asp Ser Asn Asn Asn Val Val Ala Gln Gly Lys Thr Ile Asp Asp Ser
 180 185 190

Val Glu Ala Thr Pro Lys Lys Thr Ile Glu Ser Phe Gly Gln Arg Thr
 195 200 205

Ser Ile Tyr Arg Gly Val Thr Arg His Arg Trp Thr Gly Arg Tyr Glu

210

215

220

Ala His Leu Trp Asp Asn Ser Cys Lys Arg Glu Gly Gln Thr Arg Lys
225 230 235 240

Gly Arg Gln Val Tyr Leu Gly Gly Tyr Asp Lys Glu Glu Lys Ala Ala
245 250 255

Arg Ala Tyr Asp Leu Ala Ala Leu Lys Tyr Trp Gly Thr Thr Thr Thr
260 265 270

Thr Asn Phe Pro Met Ser Glu Tyr Glu Lys Glu Val Glu Glu Met Lys
275 280 285

His Met Thr Arg Gln Glu Tyr Val Ala Ser Leu Arg Arg Lys Ser Ser
290 295 300

Gly Phe Ser Arg Gly Ala Ser Ile Tyr Arg Gly Val Thr Arg His His
305 310 315 320

Gln His Gly Arg Trp Gln Ala Arg Ile Gly Arg Val Ala Gly Asn Lys
325 330 335

Asp Leu Tyr Leu Gly Thr Phe Gly Thr Gln Glu Glu Ala Ala Glu Ala
340 345 350

Tyr Asp Ile Ala Ala Ile Lys Phe Arg Gly Leu Thr Ala Val Thr Asn
355 360 365

Phe Asp Met Asn Arg Tyr Asn Val Lys Ala Ile Leu Glu Ser Pro Ser
370 375 380

Leu Pro Ile Gly Ser Ala Ala Lys Arg Leu Lys Glu Ala Asn Arg Pro
385 390 395 400

Val Pro Ser Met Met Met Ile Ser Asn Asn Val Ser Glu Ser Glu Asn
405 410 415

Ser Ala Ser Gly Trp Gln Asn Ala Ala Val Gln His His Gln Gly Val
420 425 430

Asp Leu Ser Leu Leu His Gln His Gln Glu Arg Tyr Asn Gly Tyr Tyr
435 440 445

Tyr Asn Gly Gly Asn Leu Ser Ser Glu Ser Ala Arg Ala Cys Phe Lys
450 455 460

Gln Glu Asp Asp Gln His His Phe Leu Ser Asn Thr Gln Ser Leu Met
465 470 475 480

Thr Asn Ile Asp His Gln Ser Ser Val Ser Asp Asp Ser Val Thr Val
485 490 495

Cys Gly Asn Val Val Gly Tyr Gly Gly Tyr Gln Gly Phe Ala Ala Pro
500 505 510

Val Asn Cys Asp Ala Tyr Ala Ala Ser Glu Phe Asp Tyr Asn Ala Arg
515 520 525

Asn His Tyr Tyr Phe Ala Gln Gln Gln Gln Thr Gln Gln Ser Pro Gly
530 535 540

Gly Asp Phe Pro Ala Ala Met Thr Asn Asn Val Gly Ser Asn Met Tyr
545 550 555 560

Tyr His Gly Glu Gly Gly Glu Val Ala Pro Thr Phe Thr Val Trp
565 570 575

Asn Asp Asn

<210> 3
<211> 2011
<212> DNA
<213> Brassica napus

<400> 3
ttcttctttt acccttttacc aagaactcgt tagatcattt tctgaactcg atgaataata 60
actgggttagg cttttctctc tctccttatg aacaaaaatca ccatcgtaag gacgtctgct 120
cttccaccac cacaaccgcc gtatagtctg ccggagagta ctgtttacag ccgacccgctg 180
ctctccgatga gtcttcagcc atccaaacat cgtttctctc tccctttggt gtgcgtctcg 240
atgctttcac cagagacaac aatagtcact ccggagattg ggacatcaat ggttagtgcat 300
gtaataacat ccacaatgat gagcaagatg gaccaaaact tgagaatttc cttggccgca 360
ccaccacgat ttacaacacc aacgaaaaacg ttggagatat cgtatggaat ggggtgttatg 420
gaggaggaga cgggtggtgt ggctcactag gactttcgat gataaagaca tggctgagaa 480
atcaaccogt ggataatggt gataatcaag aaaatggcaa tggtgcaaaa ggctgtccc 540
tctcaatgaa ctcatctact tcttgtgata acaacaacta cagcagtaac aacctgttg 600
cccaagggaa gactattgat gatagcgttg aagctacacc gaagaaaact attgagagtt 660
ttggacagag gacgtctata taccgcggtg ttacaaggca tccgtgggca ggaagatatg 720
aggcacattt atgggataat agttgtaaac gagaaggcca aacgcgcaaa ggaagacaag 780
tttatttggg aggttatgac aaagaagaaa aagcagctag ggcttatgat tttagccgac 840
tcagagtattg gggaaccacc actactacta acttccocat gacgcaatat ggaaagaga 900
tagaagagat gaagacacatg acaaggccaag agtatgttgc ctactctgc aggaagaa 960
gtggtttctc tcgtgttgca tcgatttacc gtggagtaac aagacatcac caacatggaa 1020
gatggcaagc taggatagga agagtgcgcg gtaacaaaaga cctctacttg ggaactttg 1080
gcacacaaga agaagctgca gaggcatacg acattgcgpc catcaaattc agaggattaa 1140
ccgcagtgac taacttcgac atgaacagat acaacgttaa agcaatctctc gaaagcccta 1200
gtcttctcat tggtagcgcc gcaaaaacgtc tcaaggagcg taacgctccg gtccaagta 1260
tgatgatgat cagtaataac gtttcagaga gtgagaataa tgctagcgtg tggcaaaacg 1320
ctgcgggtca gcatcatcag ggagtatagt tgagcttatt gcagcaacat caagagaggt 1380
acaatgggta ttattacaat ggaggaaaact tgtcttcgga gagtgctagg gcttgtttca 1440
aacaagagga tgatcaacac catttcttga gcaacacgca gagcctcatg actaatatcg 1500
atcatcaaa gtctgtttca gatgattcgg ttactgtttg tggaaaatgtt gttggttatg 1560
gtgggttatca aggatttgca gccccggtta actcgcgatg ctacgctcgt agtgagtttg 1620
actataacgc aagaaaacat tattactttg ctacagagca gcagaccagc ctactcgccag 1680
gaggagattt tcccgcggca atgacgaata atgttggctc taatatgtat taccatgggg 1740
aaggtgggtg agaagttgct ccaacattta cagtttggaa cgacaattag aaataatagt 1800
taaagatctt tagtttatgt cgttgtgtgt tgggtgttga cagtttgata ctttgattat 1860
gttttttttt ctctttttca ttttgtgtgt tagtttctta agacttattt tttgtttcca 1920
ttagtgtgat aaattttcgg acttaagggt cactctgtgt ctgactctg tctaatacag 1980
aaaagttttc ataaaaaaa aaaaaaaaaa a 2011

<210> 4
 <211> 579
 <212> PRT
 <213> Brassica napus

<400> 4
 Met Asn Asn Asn Trp Leu Gly Phe Ser Leu Ser Pro Tyr Glu Gln Asn
 1 5 10 15
 His His Arg Lys Asp Val Tyr Ser Ser Thr Thr Thr Thr Val Val Asp
 20 25 30
 Val Ala Gly Glu Tyr Cys Tyr Asp Pro Thr Ala Ala Ser Asp Glu Ser
 35 40 45
 Ser Ala Ile Gln Thr Ser Phe Pro Ser Pro Phe Gly Val Val Val Asp
 50 55 60
 Ala Phe Thr Arg Asp Asn Asn Ser His Ser Arg Asp Trp Asp Ile Asn
 65 70 75 80
 Gly Cys Ala Cys Asn Asn Ile His Asn Asp Glu Gln Asp Gly Pro Lys
 85 90 95
 Leu Glu Asn Phe Leu Gly Arg Thr Thr Thr Ile Tyr Asn Thr Asn Glu
 100 105 110
 Asn Val Gly Asp Gly Ser Gly Ser Gly Cys Tyr Gly Gly Asp Gly
 115 120 125
 Gly Gly Gly Ser Leu Gly Leu Ser Met Ile Lys Thr Trp Leu Arg Asn
 130 135 140
 Gln Pro Val Asp Asn Val Asp Asn Gln Glu Asn Gly Asn Ala Ala Lys
 145 150 155 160
 Gly Leu Ser Leu Ser Met Asn Ser Ser Thr Ser Cys Asp Asn Asn Asn
 165 170 175
 Asp Ser Asn Asn Asn Val Val Ala Gln Gly Lys Thr Ile Asp Asp Ser
 180 185 190
 Val Glu Ala Thr Pro Lys Lys Thr Ile Glu Ser Phe Gly Gln Arg Thr
 195 200 205
 Ser Ile Tyr Arg Gly Val Thr Arg His Arg Trp Thr Gly Arg Tyr Glu
 210 215 220
 Ala His Leu Trp Asp Asn Ser Cys Lys Arg Glu Gly Gln Thr Arg Lys
 225 230 235 240
 Gly Arg Gln Val Tyr Leu Gly Gly Tyr Asp Lys Glu Glu Lys Ala Ala
 245 250 255

Arg	Ala	Tyr	Asp	Leu	Ala	Ala	Leu	Lys	Tyr	Trp	Gly	Thr	Thr	Thr	Thr
			260					265					270		
Thr	Asn	Phe	Pro	Met	Ser	Glu	Tyr	Glu	Lys	Glu	Val	Glu	Glu	Met	Lys
		275					280					285			
His	Met	Thr	Arg	Gln	Glu	Tyr	Val	Ala	Ser	Leu	Arg	Arg	Lys	Ser	Ser
		290				295					300				
Gly	Phe	Ser	Arg	Gly	Ala	Ser	Ile	Tyr	Arg	Gly	Val	Thr	Arg	His	His
					310					315				320	
Gln	His	Gly	Arg	Trp	Gln	Ala	Arg	Ile	Gly	Arg	Val	Ala	Gly	Asn	Lys
				325					330					335	
Asp	Leu	Tyr	Leu	Gly	Thr	Phe	Gly	Thr	Gln	Glu	Glu	Ala	Ala	Glu	Ala
			340					345					350		
Tyr	Asp	Ile	Ala	Ala	Ile	Lys	Phe	Arg	Gly	Leu	Thr	Ala	Val	Thr	Asn
		355				360						365			
Phe	Asp	Met	Asn	Arg	Tyr	Asn	Val	Lys	Ala	Ile	Leu	Glu	Ser	Pro	Ser
		370				375					380				
Leu	Pro	Ile	Gly	Ser	Ala	Ala	Lys	Arg	Leu	Lys	Glu	Ala	Asn	Arg	Pro
					390					395				400	
Val	Pro	Ser	Met	Met	Met	Ile	Ser	Asn	Asn	Val	Ser	Glu	Ser	Glu	Asn
				405					410					415	
Ser	Ala	Ser	Gly	Trp	Gln	Asn	Ala	Ala	Val	Gln	His	His	Gln	Gly	Val
			420					425				430			
Asp	Leu	Ser	Leu	Leu	His	Gln	His	Gln	Glu	Arg	Tyr	Asn	Gly	Tyr	Tyr
		435				440						445			
Tyr	Asn	Gly	Gly	Asn	Leu	Ser	Ser	Glu	Ser	Ala	Arg	Ala	Cys	Phe	Lys
		450				455					460				
Gln	Glu	Asp	Asp	Gln	His	His	Phe	Leu	Ser	Asn	Thr	Gln	Ser	Leu	Met
				470						475				480	
Thr	Asn	Ile	Asp	His	Gln	Ser	Ser	Val	Ser	Asp	Asp	Ser	Val	Thr	Val
				485				490						495	
Cys	Gly	Asn	Val	Val	Gly	Tyr	Gly	Gly	Tyr	Gln	Gly	Phe	Ala	Ala	Pro
			500					505					510		
Val	Asn	Cys	Asp	Ala	Tyr	Ala	Ala	Ser	Glu	Phe	Asp	Tyr	Asn	Ala	Arg
			515				520					525			
Asn	His	Tyr	Tyr	Phe	Ala	Gln	Gln	Gln	Gln	Thr	Gln	Gln	Ser	Pro	Gly
		530				535						540			
Gly	Asp	Phe	Pro	Ala	Ala	Met	Thr	Asn	Asn	Val	Gly	Ser	Asn	Met	Tyr
					550					555				560	

Tyr His Gly Glu Gly Gly Gly Glu Val Ala Pro Thr Phe Thr Val Trp
565 570 575

Asn Asp Asn

<210> 5
<211> 4873
<212> DNA
<213> Brassica napus

<220>
<221> intron
<222> (1846)..(2298)

<220>
<221> intron
<222> (2720)..(2952)

<220>
<221> intron
<222> (3036)..(3160)

<220>
<221> intron
<222> (3170)..(3314)

<220>
<221> intron
<222> (3404)..(3553)

<220>
<221> intron
<222> (3628)..(3797)

<220>
<221> intron
<222> (3849)..(3961)

<220>
<221> intron
<222> (4039)..(4148)

<220>
<221> misc_feature
<222> (1620)..(1622)
<223> start codon

<220>
<221> misc_feature
<222> (4856)..(4858)
<223> stop codon

<400> 5

8

```

cgcaactcaag tattggggaa ccaccactac tactaacttc cccgtaagtc aatcaatggt 3420
gtacaagatt tcataactta gaaccaatatt tattcttttt ttataagatg ctattatctt 3480
attatttaatt gccatgttta tatcgttaca ttattaccaa taaaaagtag ttttggtttg 3540
atataatatg tagatgagcg aatatgaaaa agaggtagaa gagatgaagc acatgacaag 3600
gcaagagtat gttgcctcac tgogcaggta tataatggaa cttctgatat tattgcatat 3660
ggcatctatt attatacatg tatattagta ttttatatat agaaccctac acgtcacgt 3720
ttatatttaa aaatatgtcc gtattcacgt cagattatca gcatacacct atatataata 3780
gacattaaaa tatgcaggaa aagttagtgg ttctctcgtg gtgcctcgat ttatcgtgga 3840
gtaccaagggt attcatacag agagaacgaa tcttattttg ttacgtacat atatatataa 3900
aaatataatt ataagatatt acattttata ttatgaatat ttcttctaatt ggggtccaaaa 3960
gcatacccca acatggaaga tggcaagcta ggataggaaag agtgcgggtg aacaaagacc 4020
tctacttggg aacttttgggt acgttttagtc ttctcttact aaacttccaa atcaaatcta 4080
taacaaaaga tatcaactaa aaactacaac atatatctaa gtaagetgta catatattat 4140
atatgaaggc acacaagaag aagctgcaga ggcatacgac attgcggcca tcaaattcag 4200
aggattaacc gcagtgacta acttcgacat gaacagatac aacgttaaag caatcctcga 4260
aagccctagt cttcctattg gtacgcgcgcg aaaaacgtctc aaggaggcta accgtccggt 4320
tccaagtatg atgatgatca gtaataacgt ttcagagagt gagaatagtg ctacgcggtg 4380
gcaaaaacgct gcggttcagc atcatcaggg agtagatttg agcttattgc accaactatca 4440
agagagggtac aatgggttatt attacaatgg aggaaaacttg tcttcggaga gtgctagggc 4500
ttgtttcaaa caagaggatg atcaacaacca ttctcttgagc aacacgcaga gcctcatgac 4560
taatatcgat catcaaaagt ctgttttcgga tgattcgggt actgtttgtg gaaatgttg 4620
tggttatggt ggttatcaag gatttgcagc cccggttaac tgcgatgcct acgctgctag 4680
tgagtttgat tataacgcga gaaaccatta ttactttgct cagcagcagc agaccagca 4740
gtcgccaggt ggagattttc ccgcggcaat gacgaataat gttggctcta atatgtatta 4800
ccatggggaa ggtgtgtggg aagttgctcc aacatttaca gtttggaacg acaattagaa 4860
aaaaatagta aag
4873

```

```

<210> 6
<211> 5151
<212> DNA
<213> Arabidopsis thaliana

```

```

<220>
<221> intron
<222> (2249)..(2578)

<220>
<221> intron
<222> (2994)..(3220)

```

```

<220>
<221> intron
<222> (3304)..(3420)

```

```

<220>
<221> intron
<222> (3429)..(3521)

```

```

<220>
<221> intron
<222> (3611)..(3770)

```

```

<220>
<221> intron
<222> (3845)..(3969)

```

<220>
 <221> intron
 <222> (4020)..(4151)

<220>
 <221> intron
 <222> (4229)..(4310)

<220>
 <221> misc_feature
 <222> (2026)..(2028)
 <223> start codon

<220>
 <221> misc_feature
 <222> (5033)..(5035)
 <223> stop codon

<400> 6
 ttctaaactc atccatctga ttttaataac agttttttct tctttttctt ttgttgtttt 60
 ttaccacttt tcttttcttt tctcattttc tactttactc cagatttttc attttcttat 120
 ttttggtcac acgctcttgt cagttgtaga tatcttcacg tacaggtgtt tctttttatt 180
 ttccagatgga atctcaatct acagggtgtt ctcacttcaa taaattacgg cccccaaaaa 240
 atttagtttt tgtattttaca agaaacatag cataaataga tacatatggt ttggaagtac 300
 tgttttttac acaaaaacttt gattataaaa cctcagcgtg tctttcgtat ttgaatttta 360
 aacgcatgca atgaagtcac tcgtgaatga tatataaata gtttgtttat ttgttatata 420
 tcgtcccgcc ccggatcaaaa acctaaagta agtgaataaa attttctttt gttagagataa 480
 gaaaatttgt accgcgtatc gaaaatgtaa aacctatttt aatttctaga tctactaatt 540
 gggtttgagg tattgaaata attgggtacc aaagggtttg ggtactatat ataaaaagca 600
 gataagaaca aattgttagg aaaaaataat atgattttgt aggtaccgag gcaattctag 660
 aacgtgtgtt ggtgtgtgtg tagatattgc aggcataata atggaagaag tgaatttata 720
 ttcaaattaa ataggaagac gagaatccat tgaatcatat cttaccagtc caaacttttt 780
 ttaagtatat aaacttttga aagagtataa acccatgcac atgcccactt tcgtctcatt 840
 gatccatgtg tataccctat agtttctctc ctaatttact taattccctt aaactatttt 900
 ttaatttgat acaattagtc ggataagctc aaactacttt actatttggt cttagcatgt 960
 acagttacata tctagcatct gaaccctact agccatccac atcttatgta cataaattag 1020
 actgttttaa gtaacttttt actttcgttt acaattgttt ttgaaaattt taggcggttt 1080
 ttacttggtt gaactgtagc cactaaagca ctaagacttc aaaattcaaa taggaaaatc 1140
 tatactttta caatattctt gcatgtcaaa ttatttttaa cgtggttata cattttgctt 1200
 aagattttag gtacattcat aataacaaac ataaaaattt tctatatata gtaggtttag 1260
 tgaagttaact atatgagata gttcatcgca ttgatccagt ctgatgcgaa tccatattcc 1320
 tatattctagt tgaacatagt tttcgtggaa gacaggaacc atctctttag ccgcgaacttc 1380
 aaaaatatac aaaaacacgaa accatgaatt ttttgagttt gttaaaaaat actaaaaagt 1440
 acgagtttgc gtttggaaaa aatgccaaac taaatcgctg gctogtgcac tacgtttcac 1500
 catacaatag tctctaaagc acacagcatc attggtotta aatcgacaac gagtgagttt 1560
 ttggactttt accatttggt cctcgacatg tttaaccatt ttgtctattt acatttaaca 1620
 ttttatacgc atgaagagag agagacagaa agcagagatt tgaattggtt tttagctgat 1680
 taattaaagt gtcatacaaaa caaatttgga ttacagattt atccagttga aacgacatta 1740
 ctaccctact ccttcaaaac gaccaatata tctccacatt tctccagtaa atattttttc 1800
 ttcttgaatt taattgcaaa attctctaaa tgcgcataat atgtcgcttc ggaagaaatg 1860
 aacattatat ttttgacttt tctctctctt cttctctctt tctctctatt taacacaaaa 1920
 accctttttt tctctctctt catgcatgaa ccctaactaa gttctttttc ctattctctt 1980
 tctctcatct tctcaacagg gtatgttaga tattatatga actcgatgaa taactgggta 2040
 ggcttctctc tctctctcct tgatcaaaat catcaccgta cggatgttga cctctccacc 2100
 accagaacccg ccgtagatgt tgccggagggt tactgttttg atctggccgc tccctccgat 2160

gaatctcttg	cogttcaaac	atcttttctt	tcctcttgcg	gtgtcaccoct	cgagcgttcc	220
acagagagaca	ataaatgctca	ctcccgaggt	ttgtgtttta	aaaatatttta	ttttattcttt	2280
gtttttttgtg	ttttttccoc	ttcttccagt	cgatagaaca	agaaacagaa	ctcacgcagc	2340
tagccctatt	ttgttttttc	attgttttate	gatttcaatc	ctttttgaga	tttccatgat	2400
tgggttttag	tgttttgttca	catgatocaa	ttctcatgaat	ttaaactgtg	taaaacatga	2460
aactagacat	ttattttgtta	coccttttate	ctttataaaat	gaaaatttoca	tttcgtatat	2520
ttatagatct	tgatgaatca	aaccocaaagt	tggggagctt	tttgttttttt	ttctatagat	2580
tgggacatca	atggtgtgtc	atgtcaaatc	ttaaccaata	acgacaacaaa	tgtagcaaaa	2640
cttgagaatt	ttctcggcgc	caccocaaag	atttacaata	ccaacagagac	cgtttgtgat	2700
ggaaatggcg	attgtggagg	agagacggct	gggtgtggcg	gctcactagc	ttcttcgatg	2760
ataaaaacat	ggctggagtta	ttctcgggtt	gcatactgct	atcatcaaga	caatggttaac	2820
gggtgcagag	ctgtcttccct	ctctatgaat	ttctctacta	gtgatagaoa	caactacaac	2880
acaacatgat	tgctcgtcca	agagaagaat	attctgttgt	tcgtagaaac	tacacagaa	2940
aaaactattg	agagtttttg	acaaaggagc	ttctatatcc	gcggtgtttac	aaggtttaat	3000
tcattgtatc	atgtatatgt	ttttgtgtgt	taaaattgtg	ttttcttggt	attgtttggg	3060
acattctaat	ggttcgggtg	agagagagct	caacggaaat	ttctctcaatg	tatatataag	3120
agaaacttat	attatgtgtac	attgggtttt	atatacaata	ctcgtctata	tatatgtgat	3180
gcctctgtac	atagtatata	atgtttgaat	ttaatgtcag	gcacoggtgc	acaggttagat	3240
acaggcgaca	tttatgggac	aatagtgtga	aaagagaagg	ccagactcgc	aaagggaagc	3300
aagtgtaact	atatataaag	ctaaattttt	aattttcatt	taccatttat	tttccaaacta	3360
atttaggttt	tcttttcaag	tgtttcaatc	aaaattgcac	ctgatggctc	tcttttcaag	3420
ttatctgggt	aagattcttga	tttttaagta	taaaattata	atagatgact	attaaatcta	3480
ttctaaagaa	ataatcaattg	ttgtgtatc	tgatctocaa	ggaggtttatg	acaaagaaga	3540
aaaagcagct	agggctttacg	attatgcgcg	ataaagtata	tggggaccoca	ccactactac	3600
taacttccoc	tgctgttaac	taatacaata	ttctatcata	atgtctcaac	tttcaaacaa	3660
ttttagtctg	aataatgccca	atctcttaaa	ctagtattat	cttactatta	actgtcatgt	3720
ttatatgttt	acaaataaaaa	ttagtaattt	tggtttgata	taatatattcag	tttgatgaa	3780
atgagaaagt	ggtagaagag	atgaaacaca	tgcagcgaga	agagatgtgt	gctcctctgc	3840
cgaggtacag	aatgaaaact	ttgaatttat	tgcaattttg	aaacccatca	cgatatattt	3900
tttataaata	tactgttaaca	tgtaaaatat	ctatttttgc	aaagataata	gaaacatgta	3960
aatatgcagg	aaaagtagtg	gtttctctct	tggtgtcatg	atttatcgag	gagtaacaag	4020
gtacgtataa	ttcatctaga	ttatggaaga	atacattgtc	ttcatatttt	tttttgtgtg	4080
atcacataata	tttgtcatat	ataactataa	attctaattg	attaatattt	cttttaaaat	4140
ggtttccaaa	tgcatcacca	acatggaagt	tggcaagcta	ggatcggaag	agtcgcggct	4200
acaaagacc	ttctatttgg	cttcttccgt	acatttttca	aaataattct	tataactata	4260
gatattaaat	atacacaat	atatctaagt	gaatcataca	aattatgtag	gcacacagga	4320
agaggtcgt	gagggcttat	actctggaac	catataaatt	agaggtataa	ggcgagtgac	4380
taacttcgac	atgacagat	acaatgttta	agcaattctc	gagagccoga	gctactacct	4440
ttgtagtctt	gcgaaacgtc	ctcaaggact	taacaatcgc	tgtccagatc	tgatgtatgt	4500
tgataacggt	tcagagagtg	caaataaagt	tgcgggttgc	caaaaacatg	cttttcagca	4560
tcactcaggga	atggatttga	gcttattgca	gcaacagcag	gagaggtacg	ttggttatta	4620
caatggaggga	aactgtgtca	ccgagagatc	tagggtttgt	gttatacaac	agggagaaca	4680
acaaacactc	ttfgaaaaact	gcgcgagta	catgactaat	gttgatcac	atagctcgac	4740
ctctgatgat	ctgttgtaacc	tttttgaaaa	tggtgttagt	tatgtgggtt	atacgaagtt	4800
cgcaatccct	tggtgaaact	cggttataaa	gcacotctct	actgctgcgt	agattgctta	4860
ccgcgcaaga	aatcaattat	actatgtcca	gcattcagca	caacacgaga	ttcagcagtc	4920
gcgcgggaga	gattttccgc	tgggcgattc	gaataaacat	agctctacaa	tgactattca	4980
cggggaaggt	ggtggagaag	ggogctcaac	gttttcagtt	tggaaacaca	actagaaaaa	5040
taagtataaag	atctttttagt	tgttttgctt	gtatgttcgc	aacagtttga	ttctgttttt	5100
ctttttctct	tttttggqta	atttttttat	aatcttttcc	taagtttcoa	t	5150

```
<210> 7
<211> 581
<212> PRT
<213> Arabidopsis thaliana
```


<400> 7

Met Asn Asn Trp Leu Gly Phe Ser Leu Ser Pro His Asp Gln Asn His
1 5 10 15

His Arg Thr Asp Val Asp Ser Ser Thr Thr Arg Thr Ala Val Asp Val
20 25 30

Ala Gly Gly Tyr Cys Phe Asp Leu Ala Ala Pro Ser Asp Glu Ser Ser
35 40 45

Ala Val Gln Thr Ser Phe Leu Ser Pro Phe Gly Val Thr Leu Glu Ala
50 55 60

Phe Thr Arg Asp Asn Asn Ser His Ser Arg Asp Trp Asp Ile Asn Gly
65 70 75 80

Gly Ala Cys Asn Thr Leu Thr Asn Asn Glu Gln Asn Gly Pro Lys Leu
85 90 95

Glu Asn Phe Leu Gly Arg Thr Thr Thr Ile Tyr Asn Thr Asn Glu Thr
100 105 110

Val Val Asp Gly Asn Gly Asp Cys Gly Gly Gly Asp Gly Gly Gly Gly
115 120 125

Gly Ser Leu Gly Leu Ser Met Ile Lys Thr Trp Leu Ser Asn His Ser
130 135 140

Val Ala Asn Ala Asn His Gln Asp Asn Gly Asn Gly Ala Arg Gly Leu
145 150 155 160

Ser Leu Ser Met Asn Ser Ser Thr Ser Asp Ser Asn Asn Tyr Asn Asn
165 170 175

Asn Asp Asp Val Val Gln Glu Lys Thr Ile Val Asp Val Val Glu Thr
180 185 190

Thr Pro Lys Lys Thr Ile Glu Ser Phe Gly Gln Arg Thr Ser Ile Tyr
195 200 205

Arg Gly Val Thr Arg His Arg Trp Thr Gly Arg Tyr Glu Ala His Leu
210 215 220

Trp Asp Asn Ser Cys Lys Arg Glu Gly Gln Thr Arg Lys Gly Arg Gln
225 230 235 240

Val Tyr Leu Gly Gly Tyr Asp Lys Glu Glu Lys Ala Ala Arg Ala Tyr
245 250 255

Asp Leu Ala Ala Leu Lys Tyr Trp Gly Pro Thr Thr Thr Thr Asn Phe
260 265 270

Pro Leu Ser Glu Tyr Glu Lys Glu Val Glu Glu Met Lys His Met Thr
275 280 285

Arg Gln Glu Tyr Val Ala Ser Leu Arg Arg Lys Ser Ser Gly Phe Ser
290 295 300

Arg Gly Ala Ser Ile Tyr Arg Gly Val Thr Arg His His Gln His Gly
305 310 315 320

Arg Trp Gln Ala Arg Ile Gly Arg Val Ala Gly Asn Lys Asp Leu Tyr
325 330 335

Leu Gly Thr Phe Gly Thr Gln Glu Glu Ala Ala Glu Ala Tyr Asp Ile
340 345 350

Ala Ala Ile Lys Phe Arg Gly Leu Ser Ala Val Thr Asn Phe Asp Met
355 360 365

Asn Arg Tyr Asn Val Lys Ala Ile Leu Glu Ser Pro Ser Leu Pro Ile
370 375 380

Gly Ser Ser Ala Lys Arg Leu Lys Asp Val Asn Asn Pro Val Pro Ala
385 390 395 400

Met Met Ile Ser Asn Asn Val Ser Glu Ser Ala Asn Asn Val Ser Gly
405 410 415

Trp Gln Asn Thr Ala Phe Gln His His Gln Gly Met Asp Leu Ser Leu
420 425 430

Leu Gln Gln Gln Gln Glu Arg Tyr Val Gly Tyr Tyr Asn Gly Gly Asn
435 440 445

Leu Ser Thr Glu Ser Thr Arg Val Cys Phe Lys Gln Glu Glu Glu Gln
450 455 460

Gln His Phe Leu Arg Asn Ser Pro Ser His Met Thr Asn Val Asp His
465 470 475 480

His Ser Ser Thr Ser Asp Asp Ser Val Thr Val Cys Gly Asn Val Val
485 490 495

Ser Tyr Gly Gly Tyr Gln Gly Phe Ala Ile Pro Val Gly Thr Ser Val
500 505 510

Asn Tyr Asp Pro Phe Thr Ala Ala Glu Ile Ala Tyr Asn Ala Arg Asn
515 520 525

His Tyr Tyr Tyr Ala Gln His Gln Gln Gln Gln Gln Ile Gln Gln Ser
530 535 540

Pro Gly Gly Asp Phe Pro Val Ala Ile Ser Asn Asn His Ser Ser Asn
545 550 555 560

Met Tyr Phe His Gly Glu Gly Gly Gly Glu Gly Ala Pro Thr Phe Ser
565 570 575

Val Trp Asn Asp Thr
580

```

<210> 8
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 8
gaggcagcgg tcggatcgta acagtactct
30

<210> 9
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 9
cataaggaga gagagaaaag cctaaccagt
30

<210> 10
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 10
accaagaact cgtagatc
19

<210> 11
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 11
aacgcatata actaaagatc
20

<210> 12
<211> 26
<212> DNA
<213> Artificial Sequence

<220>

```

<223> Description of Artificial Sequence: Primer

<400> 12

ccatggatcc agagacgaag cgaaac

26

<210> 13

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 13

actccatgga taataactgg ttaggc

26

<210> 14

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 14

aaattctcaa gctttggtcc atcttg

26

WO 00/75330

1/13

SEQUENCE LISTING

- <110> Plant Research International
 Her Majesty the Queen in Right of Canada as Represented by the
 Minister of Agriculture and Agri-Food Canada
- <120> Use of the BNM3 Transcriptional Activator to Control
 Plant Embryogenesis and Regeneration Processes
- <130> 08-887547W0
- <140>
 <141>
- <150> EP 99201745.9-2106
 <151> 1999-06-02
- <160> 14
- <170> PatentIn Ver. 2.1
- <210> 1
 <211> 2014
 <212> DNA
 <213> Brassica napus
- <400> 1
 gttcatctct tcttctttaag accaaaaacct ttttctctct ctctctcatgc atgaacctta 60
 actaagttct tcttctttta cctttttacca agaactcgtt agatcactct ctgaactcaa 120
 tgaataataa ctgggttaggc ttttctctct ctctcttatga acaaaaatcac catcgtaagg 180
 acgtctactc ttccaccacc acaaccgtcg tagatgtcgc cggagagtag tggtacagcat 240
 cgaccgtcgc ctccgatagc tcttcagcca tccaaacatc gtttctctct cctctttggg 300
 tctgtctcga tgcctttcacc agagacaaca atagtcactc cggagattgg gacatcaatg 360
 gttgtgcacg caataacatc cacaaacgat agcaagatgg accaaagctt gagaatttcc 420
 ttggccgcac caccacgatt tacaacacca acgaaaaagt tggagatgga agtggaagtg 480
 gctgttatgg aggaggagac ggtgggtggg gctcactagg actttcgtatg ataaagacat 540
 ggctgagaaa tcaaccgtgt gataatgttg ataatacaga aaatggcaat gctgcaaaaag 600
 gcctgtccct ctcaatgaac tcatctactt cttgtgataa caacaacgac agcaataaca 660
 acgtttttgc ccaagggaag actatttgat atagcgttga agctacaccg aagaaaaacta 720
 ttgagagttt tggacagagg acgtctatat accgcgtgtg tacaaggcat cgggtggacag 780
 gaagatatga ggcacattta tgggataata gttgtaaaaa agaaggccaa acgcggcaag 840
 gaagacaagt ttatttggga ggttatgaca aagaagaaaa agcagctcgg gcttatgatt 900
 tagccgcact caagatttgg ggaacaccca ctactactaa ctctcccatg agcgaatatg 960
 aaaaaagagt agaaagatg aagcacatga caaggcaaga gttgtgtgct tcaactggca 1020
 aacatcagat atggcaagct aggtaggcat cgatttatcg tggagtaaca agacatcacc 1080
 gaacttttgg cacacaagaa gaagctgcag aggcatacga caactgcggc atcaaatcca 1200
 gaggattaac cgcagtactt aacttcgaca tgaacagata caactgttaa gcaatctcgc 1260
 aaagccctag tcttcttatt ggtagccgcg caaaacgtct caaggagctt aaccgtcccg 1320
 ttccaagtat gatgatgact agtaataacg tttcagagag tgagaaatagt gctagcgttt 1380
 ggcaaaaacgc gctggttcag catcatcagg gactatattg gacttatgtg caccaacatc 1440
 aagagaggtta caatggttat tattacaatg gaggaacctt gcttctggag agtgctaggg 1500
 ctgttttcaa acaagaggat gatcaacacc atttcttgag caacaacatg agctcatgta 1560
 ctaatatcga toatcaaatg totgtttcgg atgattcgtt tactgttttg ggaaatgttg 1620
 ttgtttatgg tggtttcaaa ggaatttcga ccccggttaa ctgcgatgcc tacgctgcta 1680
 gctgagtttga ttataaccca agaaccattt attactttgc tcagccagcag cagaccagc 1740
 agtgcgcagg tggagatttt cccgcggcaa tgacgaataa gtttggctct aatatgtatt 1800
 accatgggga aggttggttga gaagtgtctc caacatttac agtttggaac gacaattaga 1860
 aaaaatagtt aaagattctt agttatatgc gttgttgtgt ctgtgtgaac agtgtgtgat 1920
 ttgtattatg tttttttctt tctctttctt ttttctcttg ttaatttctt aagactttat 1980
 tttagttctc attagtttga taaattttca gact 2014
- <210> 2
 <211> 579

2/13

<212> PRT

<213> Brassica napus

<400> 2

```

Met Asn Asn Asn Trp 5 Leu Gly Phe Ser 10 Leu Ser Pro Tyr Glu Gln Asn
1 5 10 15
His His Arg Lys Asp Val Tyr Ser Ser Thr Thr Thr Thr Val Val Asp
20 25 30
Val Ala Gly Glu Tyr Cys Tyr Asp Pro Thr Ala Ala Ser Asp Glu Ser
35 40 45
Ser Ala Ile Gln Thr Ser Phe Pro Ser Pro Phe Gly Val Val Val Asp
50 55 60
Ala Phe Thr Arg Asp Asn Asn Ser His Ser Arg Asp Trp Asp Ile Asn
65 70 75 80
Gly Cys Ala Cys Asn Asn Ile His Asn Asp Glu Gln Asp Gly Pro Lys
85 90 95
Leu Glu Asn Phe Leu Gly Arg Thr Thr Thr Ile Tyr Asn Thr Asn Glu
100 105 110
Asn Val Gly Asp Gly Ser Gly Ser Gly Cys Tyr Gly Gly Gly Asp Gly
115 120 125
Gly Gly Gly Ser Leu Gly Leu Ser Met Ile Lys Thr Trp Leu Arg Asn
130 135 140
Gln Pro Val Asp Asn Val Asp Asn Gln Glu Asn Gly Asn Ala Ala Lys
145 150 155 160
Gly Leu Ser Leu Ser Met Asn Ser Ser Thr Ser Cys Asp Asn Asn Asn
165 170 175
Asp Ser Asn Asn Asn Val Val Ala Gln Gly Lys Thr Ile Asp Asp Ser
180 185 190
Val Glu Ala Thr Pro Lys Lys Thr Ile Glu Ser Phe Gly Gln Arg Thr
195 200 205
Ser Ile Tyr Arg Gly Val Thr Arg His Arg Trp Thr Gly Arg Tyr Glu
210 215 220
Ala His Leu Trp Asp Asn Ser Cys Lys Arg Glu Gly Gln Thr Arg Lys
225 230 235 240
Gly Arg Gln Val Tyr Leu Gly Gly Tyr Asp Lys Glu Glu Lys Ala Ala
245 250 255
Arg Ala Tyr Asp Leu Ala Ala Leu Lys Tyr Trp Gly Thr Thr Thr
260 265 270
Thr Asn Phe Pro Met Ser Glu Tyr Glu Lys Glu Val Glu Glu Met Lys
275 280 285
His Met Thr Arg Gln Glu Tyr Val Ala Ser Leu Arg Arg Lys Ser Ser
290 295 300
Gly Phe Ser Arg Gly Ala Ser Ile Tyr Arg Gly Val Thr Arg His His
305 310 315 320
Gln His Gly Arg Trp Gln Ala Arg Ile Gly Arg Val Ala Gly Asn Lys
325 330 335

```

SUBSTITUTE SHEET (RULE 26)

3/13

Asp Leu Tyr Leu Gly Thr Phe Gly Thr Gln Glu Glu Ala Ala Glu Ala
 340 345 350
 Tyr Asp Ile Ala Ala Ile Lys Phe Arg Gly Leu Thr Ala Val Thr Asn
 355 360 365
 Phe Asp Met Asn Arg Tyr Asn Val Lys Ala Ile Leu Glu Ser Pro Ser
 370 375 380
 Leu Pro Ile Gly Ser Ala Ala Lys Arg Leu Lys Glu Ala Asn Arg Pro
 385 390 395 400
 Val Pro Ser Met Met Met Ile Ser Asn Asn Val Ser Glu Ser Glu Asn
 405 410 415
 Ser Ala Ser Gly Trp Gln Asn Ala Ala Val Gln His His Gln Gly Val
 420 425 430
 Asp Leu Ser Leu Leu His Gln His Gln Glu Arg Tyr Asn Gly Tyr Tyr
 435 440 445
 Tyr Asn Gly Gly Asn Leu Ser Ser Glu Ser Ala Arg Ala Cys Phe Lys
 450 455 460
 Gln Glu Asp Asp Gln His His Phe Leu Ser Asn Thr Gln Ser Leu Met
 465 470 475 480
 Thr Asn Ile Asp His Gln Ser Ser Val Ser Asp Asp Ser Val Thr Val
 485 490 495
 Cys Gly Asn Val Val Gly Tyr Gly Gly Tyr Gln Gly Phe Ala Ala Pro
 500 505 510
 Val Asn Cys Asp Ala Tyr Ala Ala Ser Glu Phe Asp Tyr Asn Ala Arg
 515 520 525
 Asn His Tyr Tyr Phe Ala Gln Gln Gln Gln Thr Gln Gln Ser Pro Gly
 530 535 540
 Gly Asp Phe Pro Ala Ala Met Thr Asn Asn Val Gly Ser Asn Met Tyr
 545 550 555 560
 Tyr His Gly Glu Gly Gly Gly Glu Val Ala Pro Thr Phe Thr Val Trp
 565 570 575
 Asn Asp Asn

<210> 3
 <211> 2011
 <212> DNA
 <213> Brassica napus

<400> 3
 ttctttttttt acctttttacc aagaactcgt tagatcattt tctgaactcg atgaataata 60
 actgggttagg cttttctctc tctctttatg aacaaaatca ccatcgtaag gacgtctgct 120
 cttccaccac cacaaccgcc gtatagtctg cgggagagta ctgttacgat ccgaccgctg 180
 cctccgatga gtcttcagcc atccaaacat cgtttctctc tccctttggt gtcgtctctg 240
 atgctttcac cagagacaac aatagtcact cccgagattg ggacatcaat ggtagtgcac 300
 gtaataacat ccacaatgat gagcaagatg gaccaaaact tgagaatttc cttggccgca 360
 ccaccagat ctcaacacc aacgaaaaac ttggagatat cgatggaagt ggggtgttatg 420
 gagagggaga cggtcggtgt ggctcactag gactttcgat gataaagaca tggctgagaa 480
 atcaaccggt ggaataatgt gataatcaag aaaaatggca cggctgcaaaa aacctgtccc 540
 tctcaatgaa tctactact tcttgtgata acaacaacta cagcagtaac aacctgtgtg 600
 cccaagggaa gactattgat gatagcgttg aagctacacc gaagaaaact attgagagtt 660

4/13

ttggacagag gacgtctata taccgcggtg ttacaaggca tcgggtggaca ggaagatatg 720
 aggcacattt atgggataat agttgtaaac gagaaggcca aacggcgcaa ggaagacaag 780
 tttattttgg aggttatgac aaagaagaaa aagcagctag ggcttatgat ttgaccgcac 840
 tcaagtattg gggaaccacc actactacta acttccccat gagcgaaatag gagaagaaga 900
 tagaagagat gaagcacatg acaaggcaag agtatgttgc ctccacttcgc agggaaaagta 960
 gtggtttttc tcgtggtgca tcgattttatc gtggagtaac aagacatcac caacatggaa 1020
 gatggcaagc taggtatagga agagtccgcc gtaacaaaga cctctactctt ggaacttttg 1080
 gcacacaaga agaagctgca gaggcatacg acattgcggc catcaaatc agaggattaa 1140
 ccgcagatgac taacttcgac atgaacagat acaacgttaa agcaatctct gaaagcccta 1200
 gtcttcttat tggtagcgcc gcaaaacgtc tcaaggaggc taaccgtccg ttccaagta 1260
 tgatgatgat cagtataaac gtttcagaga gtgagaataa tgctagcggt tggcaaacg 1320
 ctgcggttca gcatcatcag ggagtagatt tgagcttatt gcagcaacat caagagaggt 1380
 acaatggtta ttattacaat ggaggaaact tgtcttcgga gagtgtctagg gcttggttca 1440
 aacaagagga tgatcaacac catttcttga gcaacacgca gagcctcatg actaatcatg 1500
 atcatcaag ttctgtttca gatgattcgg ttactgtttg tggaaatgtt gttggttatg 1560
 gtggttatca aggattttgca gccccggtta actcgcatgc ctacgctgct agtgagtttg 1620
 actataacgc aagaaaccat tattactttg ctacgacgca gcagaccacg catcggccag 1680
 gaggagtttt tcccgcggca atgacgaata atgttggctc taatagttag taccatgggg 1740
 aaggttggtg agaagttgct ccaacattta cagtttggaa cgacaattag aaataatagt 1800
 taaagatctt tagttatatg cgttgttggt tgggtgttga cagtttgata -cttgattat 1860
 gttttttttt ctctttttca ctttgttggt tagtttctta agacttattt tttgtttcca 1920
 ttagtggat aaatttttcg acttaagggt cacttctggt ctgacttctg tctaatacacg 1980
 aaaaagtttc ataaaaaaaa aaaaaaaaaa a 2011

<210> 4
 <211> 579
 <212> PRT
 <213> Brassica napus

<400> 4
 Met Asn Asn Asn Trp Leu Gly Phe Ser Leu Ser Pro Tyr Glu Gln Asn
 1 5 10 15
 His His Arg Lys Asp Val Tyr Ser Ser Thr Thr Thr Thr Val Val Asp
 20 25 30
 Val Ala Gly Glu Tyr Cys Tyr Asp Pro Thr Ala Ala Ser Asp Glu Ser
 35 40 45
 Ser Ala Ile Gln Thr Ser Phe Pro Ser Pro Phe Gly Val Val Val Asp
 50 55 60
 Ala Phe Thr Arg Asp Asn Asn Ser His Ser Arg Asp Trp Asp Ile Asn
 65 70 75 80
 Gly Cys Ala Cys Asn Asn Ile His Asn Asp Glu Gln Asp Gly Pro Lys
 85 90 95
 Leu Glu Asn Phe Leu Gly Arg Thr Thr Thr Ile Tyr Asn Thr Asn Glu
 100 105 110
 Asn Val Gly Asp Gly Ser Gly Ser Gly Cys Tyr Gly Gly Gly Asp Gly
 115 120 125
 Gly Gly Gly Ser Leu Gly Leu Ser Met Ile Lys Thr Trp Leu Arg Asn
 130 135 140
 Gln Pro Val Asp Asn Val Asp Asn Gln Glu Asn Gly Asn Ala Ala Lys
 145 150 155 160
 Gly Leu Ser Leu Ser Met Asn Ser Ser Thr Ser Cys Asp Asn Asn Asn
 165 170 175
 Asp Ser Asn Asn Asn Val Val Ala Gln Gly Lys Thr Ile Asp Asp Ser
 180 185 190

5/13

Val Glu Ala Thr Pro Lys Lys Thr Ile Glu Ser Phe Gly Gln Arg Thr
 195 200 205
 Ser Ile Tyr Arg Gly Val Thr Arg His Arg Trp Thr Gly Arg Tyr Glu
 210 215 220
 Ala His Leu Trp Asp Asn Ser Cys Lys Arg Glu Gly Gln Thr Arg Lys
 225 230 235 240
 Gly Arg Gln Val Tyr Leu Gly Gly Tyr Asp Lys Glu Glu Lys Ala Ala
 245 250 255
 Arg Ala Tyr Asp Leu Ala Ala Leu Lys Tyr Trp Gly Thr Thr Thr Thr
 260 265 270
 Thr Asn Phe Pro Met Ser Glu Tyr Glu Lys Glu Val Glu Glu Met Lys
 275 280 285
 His Met Thr Arg Gln Glu Tyr Val Ala Ser Leu Arg Arg Lys Ser Ser
 290 295 300
 Gly Phe Ser Arg Gly Ala Ser Ile Tyr Arg Gly Val Thr Arg His His
 305 310 315 320
 Gln His Gly Arg Trp Gln Ala Arg Ile Gly Arg Val Ala Gly Asn Lys
 325 330 335
 Asp Leu Tyr Leu Gly Thr Phe Gly Thr Gln Glu Glu Ala Ala Glu Ala
 340 345 350
 Tyr Asp Ile Ala Ala Ile Lys Phe Arg Gly Leu Thr Ala Val Thr Asn
 355 360 365
 Phe Asp Met Asn Arg Tyr Asn Val Lys Ala Ile Leu Glu Ser Pro Ser
 370 375 380
 Leu Pro Ile Gly Ser Ala Ala Lys Arg Leu Lys Glu Ala Asn Arg Pro
 385 390 395 400
 Val Pro Ser Met Met Met Ile Ser Asn Asn Val Ser Glu Ser Glu Asn
 405 410 415
 Ser Ala Ser Gly Trp Gln Asn Ala Ala Val Gln His His Gln Gly Val
 420 425 430
 Asp Leu Ser Leu Leu His Gln His Gln Glu Arg Tyr Asn Gly Tyr Tyr
 435 440 445
 Tyr Asn Gly Gly Asn Leu Ser Ser Glu Ser Ala Arg Ala Cys Phe Lys
 450 455 460
 Gln Glu Asp Asp Gln His His Phe Leu Ser Asn Thr Gln Ser Leu Met
 465 470 475 480
 Thr Asn Ile Asp His Gln Ser Ser Val Ser Asp Asp Ser Val Thr Val
 485 490 495
 Cys Gly Asn Val Val Gly Tyr Gly Gly Tyr Gln Gly Phe Ala Ala Pro
 500 505 510
 Val Asn Cys Asp Ala Tyr Ala Ala Ser Glu Phe Asp Tyr Asn Ala Arg
 515 520 525
 Asn His Tyr Tyr Phe Ala Gln Gln Gln Gln Thr Gln Gln Ser Pro Gly
 530 535 540

6/13

Gly Asp Phe Pro Ala Ala Met Thr Asn Asn Val Gly Ser Asn Met Tyr
 545 550 555 560

Tyr His Gly Glu Gly Gly Gly Glu Val Ala Pro Thr Phe Thr Val Trp
 565 570 575

Asn Asp Asn

<210> 5
 <211> 4873
 <212> DNA
 <213> Brassica napus

<220>
 <221> intron
 <222> (1846)..(2298)

<220>
 <221> intron
 <222> (2720)..(2952)

<220>
 <221> intron
 <222> (3036)..(3160)

<220>
 <221> intron
 <222> (3170)..(3314)

<220>
 <221> intron
 <222> (3404)..(3553)

<220>
 <221> intron
 <222> (3628)..(3797)

<220>
 <221> intron
 <222> (3849)..(3961)

<220>
 <221> intron
 <222> (4039)..(4148)

<220>
 <221> misc_feature
 <222> (1620)..(1622)
 <223> start codon

<220>
 <221> misc_feature
 <222> (4856)..(4858)
 <223> stop codon

<400> 5
 attctctccac cgatttcgtta cccagtgctt gaaaaatatga tgactacgaa tcaattaaat 60
 ggagaagcttc cactgcttgt gtagggtggaa gctcaagcaa caaccggaaa cctcggcggt 120
 atcgggagatt agcatcggtta ttgccaataa ttccgcgcgc agagatgaaa cgattcaaga 180
 gaaaccctca aatagggttag ccataaaaca gtgaattagt atgatttaag agataagaag 240
 agaagtcgag ttcaagaaaa gaaatactca catctattta tctgtttac acacgcgctt 300
 tcaatctaa gcaaacgatt gaagatgaat cgtggaggag agttaatagg atttaacaca 360
 aagccattaa ccaaaccggt gcaggtcggg agacgaaccg caaaagtcac gcttagccgt 420
 cgcacgaaga ggagcgatga atttcgtttt ctgcgtcgag tcgtattagg gatagacgga 480

gccattatc gttggggcgg aaacacttct aatctcacag cccatgaaca cactaaagaa 540
 cgaaccggaa aatgtttgaa gtttaagtga acgtgcggtt tgcttatgag acacatgtca 600
 ttacgatcatg aaatgatttta tctacgtgga tcataggtgt ctctcttaagg agagagcaaa 660
 cctacactctt acataaatac atttgatca tcttaagagg tgtttaagat ttttgcataa 720
 atatttaaaaa aaaaatacaaa tttttatgta actagttttg gttacataaaa ataacattaa 780
 ataaaaattaa tccaaccaat aaaaaatac ggtattttta aattggtcaa aaataaaaaa 840
 aaaaactata atttcaccta gaattacgag aatgtcactt attttgaaac aaaaataaaa 900
 tctttaacaa tcaattaaac tgatacggat ggagtataa tctttacata taacataaat 960
 atatgtttttt cctgtaagcg tccatctctt ctttagtcac tagttcaaat accagctgca 1020
 gtaaaacatc gaattattga atttgttgta aaatatctga agcagctttg 1080
 aagcaaaacg ccaaacgcga tcgctcgctc ggctcatagg tccacacata acatgtgact 1140
 agcatattgg gtcttaattc aacagcgagt gattttggga taattatata gttctcgctg 1200
 tactcttcaat ttaacacaaa gtcactaac cttatttac atgaagagag gtttgaaagg 1260
 gcttttgatg gatttaattt aatgtattaa accaaactag aattaagaga tttagcattg 1320
 aattacattta caccaccacc ccaccattca aaccgacca tacaattctca cagttttcaa 1380
 gtaaaacacac ttttttttgt tgctctctcg gaatttttaa aatatattctg ttataataat 1440
 gcgcatgata tgcgcctctg gaagaaatga aacattatatt ctttgacttt tcttctccta 1500
 gttcatctct accaaaacct tttctctctc tttctctctc agaatctctg atgaacctca 1560
 actaaagtct acttctttta cttcttaaca agaatctctg agatcactct ctgaactcaa 1620
 tgaataataa cctgttagcg tttctctctc cctctctatga acaaaatac catcgttaag 1680
 acgtctactc tctcacacc tcaaccgtcg tagatgtcgc gtcgtctctc cctcttggtg 1740
 cgacccgtct cctcttagcg tcttcacgca tccaacactc cgtttctctc tttgttagaa 1800
 tctgtctcga tggctttcac agagacaaca atagtcaact cctgtctcga tttgttagaa 1860
 ctacttgttt ttttttgtat tgtttatttg ttttagttcc tcttcttcca atgttagaa 1920
 caaagaccaa tacacacgca cgcatactag cctattttt cctctgggct ttataataat 1980
 ttttctattt ttttgtgata tcaatgtgtg gggtttgatg tttgtttgca tatagtaata 2040
 ctaaaaataa tggcagatta acatagattt ttttaaaaga tttacataga tagaaatga 2100
 aatttgacat tctctctttt attcaatctc ataatatgat cacatacatg tgtactcttt 2160
 gatttgtata tttgtttctt acagtttgaag gagagaataa ccaaatctcc atttgtatat 2220
 tatagatcgg tggtaaaag taaatttaac aaattatgat aatatagacc attaatcttt 2280
 gattttcgg tcttatagat tgggacatca atgtgtgtgc atgcaataac attccaacg 2340
 atgagcaaga tggacaaaag cttgagaatt tcttggcgcc caccacacg atttcaaca 2400
 ccaacgaaaa cgttggagat ggaagtggaa gtggctgtta tggagagaga gacggttggt 2460
 tgggtctcact aggaatttcc atgataaaga catggctgag aatacaaccc ttgggataat 2520
 ttgataatca agaaaaatgc aatgctgcga aaggcctgtc cctctcaatg aactcatcta 2580
 cttcttgtga taacacaac gacagcaata acaacgttgt tgcaccaagg aagactatgt 2640
 atgatacggc tgaagtggat ccgaagaaaa ctattgagag ttttggcag agagactcta 2700
 tatacccggtg tctcacaagg tgccttctat ttatttaatt aaaatgtcta aaatgtcgct 2760
 tgaatttgtta tcttcttggt aaagtctggg acattgatct attggctcta tgggctata 2820
 gctaccgaat ggtctctgat atatagtatc aaagagagat attgttatta tgggctata 2880
 tagaaataa tatatacata tttatgtctg tgatgacatg taagtctcta 2940
 taaatgata aggcattcgt ggacgggaag atatgaggca cattttatggg atcaatagtt 3000
 taaaagagaa gtcgaacgcg gcaaaaggaa acaaggtata tatatatctca tttgataatt 3060
 gatcataatt tctacacaga ttaatttcta acatcaataa ggtcttttga tcaattgtca 3120
 tgtttttatc aaaaatttga cctgttggtt gttctctcag atataatca attttgttat 3180
 attataaatt ggcagaagct tgatgggcta aatctaaatt atataatca tttgtttgtg 3240
 ttttttgtga tacattctat atataatca aatagcgata cgtactacat atttttagc 3300
 tcttatcatc tcaggaggtt atgacaaaga agaaaaagca cgtagggctt atgatttagc 3360
 cgcactcaag tattggggaa ccaccactac tactaaactc ccgtaagtc aatcaatgt 3420
 gtacaagatt tcaacaacta gaaccaattt ttattctttt ttataagatg cttattctct 3480
 attatttaatt gcatgttta tatcgttaca tttttacaa taaaagatc ttttggttgt 3540
 atataaatg tatatgaaa aatagtaaaa agaggtagaa gagatgaagc acatgaacg 3600
 cgaagagtat gtgcccacac tgcgcaggta taaattggaa cttctgtatg tatgtcatat 3660
 ggcattcatt attatactg tatattagta ttttatata agaaccacat agcctcacgt 3720
 tttattttaa aaatattgct gtattacagt cagattatca gcatcacctt atataataa 3780
 gacattaaaa tatgcaggaa aagtattggt tctctctgtg tggatcactt tatctgtgga 3840
 gtaacagggt attatcacag agagaacgaa tctattttg ttactgtcat acatataata 3900
 aaaaataatt ataagatatc acattttata ttatgaatat ttactctact aaacttca 3960
 gacatcaca acatttgaag tggcaagcta ggaatggaag agtcgctggt acaaaagacc 4020
 tctactctgg aaacttttagt cgttttagtc tttcttact aaacttcaac atcaaaacta 4080
 taacaaaaag tatcaactaa aaactcaaac atatatctaa gtaagctgta catatattat 4140
 atatgaagcc acacaagata aagctgcaga ggcatacgac attgcggcca tcaaatcag 4200
 aggatcaaac cagatgacta acttcgacat gaacagatca acgcgttaag caactctcga 4260
 aagccctagg gttcttaatt gtagtaact aaaaactctc aagagggatg accgtccggt 4320
 tccaagtatc atgatgact agtatgact ttcagagagt gagaattgtc ccagcggtgt 4380
 gcaaaacgct cgggttcacg atcatcagg agcagattgt agcttatgtg acccaatca 4440

8/13

```

agagagggtac aatgggttatt attacaatgg aggaacttg tcttcggaga gtgctagggc 4500
ttgtttcaaa caagaggatg atcaacacca tttcttgagc aacacgcaga gctccatgac 4560
taaatcatgat catcaaaagt ctgtttcggg tgattcgggt actgtttgtg gaaatgttgt 4620
tggttatgggt ggttatcaag gatttgcagc ccggtttaac tgcgatgctt acgctgctag 4680
tgagtttgtat tataacgcaa gaaaccatta ttactttgct cagcagcagc agacccagca 4740
gtcggccaggt ggagattttc ccgcggaat gacgaataat gttggctcta atatgtatta 4800
ccatggggaa ggtgtggag aagtgtgccc aacattttaca gtttggaacg acaattagaa 4860
aaaatagtta aag                                     4873

```

```

<210> 6
<211> 5151
<212> DNA
<213> Arabidopsis thaliana

```

```

<220>
<221> intron
<222> (2249)..(2578)

```

```

<220>
<221> intron
<222> (2994)..(3220)

```

```

<220>
<221> intron
<222> (3304)..(3420)

```

```

<220>
<221> intron
<222> (3429)..(3521)

```

```

<220>
<221> intron
<222> (3611)..(3770)

```

```

<220>
<221> intron
<222> (3845)..(3969)

```

```

<220>
<221> intron
<222> (4020)..(4151)

```

```

<220>
<221> intron
<222> (4229)..(4310)

```

```

<220>
<221> misc_feature
<222> (2026)..(2028)
<223> start codon

```

```

<220>
<221> misc_feature
<222> (5033)..(5035)
<223> stop codon

```

```

<400> 6
tc:caaaactc atccatctga ttttaataac agttttttct tctttttctt ttgttgtttt 60
ttaccacttt ttcttttttt ttctattttc tacttacttc cagatttttc atttttctat 120
tttcgggcac acgcttttgt cagttgtaga tatcttcac tacagggtgt tctttttatt 180
ttcagatgga atctcaatct acaggtgttt ctcaattcaa taaattacgg cccccaaaaa 240
atttagtttt tgcatttaca agaaacatag cataatatga tacatatggt tttgaagta 300
tg:tttttac acaaaacttt gattataaaa cctcagccgt tctttcgtat ttgaaattta 360
aacgcacgca atgaagtcac tcgtgaatga tatataataa gtttgtttat ttgttatata 420
tcgtcccgcc ccggatcaaa acctaaagta agtgaataaa attttctttt gtgagataaa 480

```

gaaaatttgg accgcgtatc gaaaatgtaa aacctatttt aattttctaga tctactaatt 540
 gggttttgagg tattgaaata attgggtacc aaaggttttgg gttactatat ataaaaagca 600
 gaataagaaac aattgcttagg aaaaaataat atgatttttgg aggtaccagg gcaatttctag 660
 aacgctgtgtt ggtgtgtgtg tagataattgc aggcataata atggaagaag tgaattata 720
 ttacaacttaa taggaaagac gagaatccat tgaatcatat ctaccaggtc caaacttttt 780
 ttaagtatat aaactctttga aagagtataa acccattgac atgcccatct tgcctccatt 840
 gatccatgtg tataccattat ggaataagctc aaacttactt acttatttggtt aaactcattt 900
 ttaatttgat tcatgcatctt gaacccatctt agccatccac atcttattgta cataaattatg 1020
 acagtacata gtaactttttt cacttaagaca ctaagacttct tctgaaattt gaggcgcttt 1080
 actgttttaa gaaactgtagc caatcatcttt gcatgtcaca ttatttttaa cgtggttata ctttatttga 1140
 ttactgtggt taactattttt caactctttt gtaacttttca ataaaaattt tcttatatata gtagggttagt 1260
 tatactttta gtaactttttt gtaactttttt gttcatgtcga ttgatgcgaa tccatatctc 1320
 aagattttga atactttttt ttaactttttt ttccgttgaa atcttcttaga cccgcacttc 1380
 gtaagtttact atatgagata gtaactttttt accatgaaac ttgtgagttt ttttaaaaaa tctgttctga 1440
 tatattctagt tgaactttttt ttccgttgaa ttgtgagttt ttttaaaaaa gctgttctga 1500
 aaaaattatca aaaaacacga accatgaaac taaactcgtg atgtgttctta aatcgacaaac 1560
 acgagttcgc cctttaagag acacagcatc attgtgtctta ttgtccattt ttgtcttctt 1620
 catcacatgt tctctattggt cctgcacatg agagacagaa agcagagatt tgaaaagggt ttgtgactga 1680
 ttgtgacttt taattaaagt gttcatgaaac caaattggga ttacagagatt atccagttagt acagacata 1740
 ctaccctcat ccttcaaacac gaccaataca tctccacatt ttccaagatt atattttttt 1800
 ttctctgaatt taattgtcaaa atttctttaa tgcgcataat atgtgccttc ggaagaagt 1860
 aacattatatt ttgtgacttt tcttcttctt cttctctctt tcttcttctt taacttggtta 1920
 acctttttct ttgttctctt catgcatgaa ccttaactaa gttcttcttt ctattctctt 1980
 ttctctcatc atcaacagga gtatgttagaa tattatatga atctcgatgaa taacttggtta 2040
 ggtcttctct tctctctcta tgaataaaa catcacccgta cggatgttga cttctccacc 2100
 accgaacccg ccgttagatgt tgcgggaggg tactgttttt atctggccgc tccctccgat 2160
 gaattctctg ccgttcaaac atctttttct tctctttctg gtgtccacct cgaagctttc 2220
 accagagaga ataatagtca ctcccgaggt ttgtgtttta aaaaatttta ttattctttt 2280
 gtttttgtta ttttttcccc ttcttccaat catagaaaca agcagacaat gtttctttt 2340
 tagccctatt ttgttttttc attgtttatc gatttcatct ctttttgaaa ttcccatgag 2400
 tggggtttag ttgtttgttc catgatcaaa ttctcatgaat ttaaaattag taaaacatga 2460
 aactagacat ttatttttga cctttttatc cttataaaat gaaaatttcca ttctgtatat 2520
 tatagatcgg tgatgaaata aacccaaacgt tggggatcgc tttaactgaa ttgtttttt 2580
 tgggacatca atgtgtgtgc atgcaataca ttaaccaata acgaacaaaa tggaccaag 2640
 cttgagaatt tcttcggccg caccaccacg atttacaata ccaacgagac cttttagat 2700
 ggaattggcg atgtgtgaggg agggagacgg ggtgtgtggc gttcactagg ctttctgat 2760
 aaaaaaacat ggtctgagtaa tctatcggtt gctaatgcta atcatcaaga caatgttaac 2820
 gttcagcag agttgttccc ctctatgaat ttgttgtatg atgtatagcaa caactacaa 2880
 acaactgatg atgtgttcca agagaagaat attgttgtatg tctatatacc ggtgtgaaac 2940
 aaaaactatt agattatatt ttatttgtct tctatatacc ggtgtgttgc atgtgttggg 3000
 acatttcaat ggttccgtgt agagagagtg caacggaatg atactattga ttcttcttgc 3060
 agaaacatta attagtgtac atgggtttat atatacaata atactgtcaa tatgttcat 3120
 gcttctgtac atagtataa atgtttgaaat aaagtgtcag gtacgtgttg acaggttagat 3180
 accgagcaca ttatgtggac aaatgttgca aaagagaagg ccagactcgc aaaggaagac 3240
 aggttactat atataataag ctaatttttt aattttcatt taccattatt atataacta 3300
 attatggttt ttatttccag ttgttcatca aaatttgac ctgatggtct tcttttctag 3360
 ttctgtgggt aagttcttga ttttaagcta taaattataa atagatgact atataacta 3420
 ttctagcaaa aataataatt ttgtgttatc tgatcctaca ggaggttagt acaaagaaga 3480
 aaaaagacgt agggctattc atttagccgc actaaagta tgggggacca ccaactacag 3540
 raacttcccc gtatgttaat taatcaataa tatatacata aattcctaac tcttaaccaa 3600
 tttttgtctt aataattgcca atctcttaaa ctagtattat ctactacta actgtcatgt 3660
 ttactattgtt acaataaaaa ttagttaatt ttgtttgata taattattcag ttgagtgaat 3720
 atagaaaga ggttagaagg atgaagcaca ttgaatttat tgacgagcca agagtattgt 3780
 gcaggttaga aagcaaaact ttgaatttat ttgatttttag aaacccatca cgtatatatt 3840
 tatttaaaaa tatctgaatc ttgaataaat ttgaatttat ttgatttttag aaacccatca cgtatatatt 3900
 aatatcagg gtattcgtgc gtttctctctg atactagtgt atactagtgt atactagtgt 3960
 gtacgtataa tccatctaga tatggaaacga atcaactta atcaactta atcaactta 4020
 atactataa atgtctatca acatgggaag ttggaaagca atcaactta atcaactta 4080
 gtttccaaa gttcatccca acattctggg aacttctggg gaatacaca atattatgat 4140
 acaaaagacc gttactctgg aacttctggg gaatacaca atattatgat atattatgat 4200
 gatattaaat atacacaaat atattctagg caatacaca atattatgat atattatgat 4260
 agaggtcgtc gaggctttat acattgcgc caatacaca atattatgat atattatgat 4320
 caactcgcac atgaacagat acaattgtta agcaatctct gagagccga ggtcactaat 4380
 caactcgcac atgaacagat acaattgtta agcaatctct gagagccga ggtcactaat 4440

10/13

```

tggtagttct gcgaaacgtc tcaaggacgt taacaatccg gttccagcta tgaatgattag 4500
taataacggt tcagagagtg caaataatgt tagcgggttg caaaacactg cgttttcagca 4560
tcatacaggga atggatttga gcttatttga gcaacagcag gagaggtagc ttggttatta 4620
caatggaggga aacttgtcta ccgagagtac taggggttgt ttcaaacaaag aggaggaaaca 4680
acaacacttc ttgagaaact cgcgagtgca catgactaat gttgatcatc atagctcgac 4740
ctctgatgat tctgttaccc ttgtggaaa tgttgttagt tatgggtggtt atcaaggatt 4800
cgcaaacctt gttggaacat cgtttaatta cgtccctttt actgctgctg agatttgcata 4860
caacgcaaga aatcattatt actatgctca gcatcagcaa caacagcaga ttcagcagtc 4920
gccgggagga gattttccgg tggcgatttc gaataaccat agctctaaca tgtactttca 4980
cggggaaggt ggtggaagaag gggctccaac gttttcagtt tgggaacgaca cttagaaaaa 5040
taagtaaaag atcttttagt tgtttgcttt gtatgttgcg aacagtttga tctctgtttt 5100
cttttccctt tttttgggta attttcttat aacttttttc atagttttca t 5151

```

<210> 7

<211> 581

<212> PRT

<213> Arabidopsis thaliana

<400> 7

```

Met Asn Asn Trp Leu Gly Phe Ser Leu Ser Pro His Asp Gln Asn His
  1          5          10          15
His Arg Thr Asp Val Asp Ser Ser Thr Thr Arg Thr Ala Val Asp Val
          20          25          30
Ala Gly Gly Tyr Cys Phe Asp Leu Ala Ala Pro Ser Asp Glu Ser Ser
  35          40          45
Ala Val Gln Thr Ser Phe Leu Ser Pro Phe Gly Val Thr Leu Glu Ala
  50          55          60
Phe Thr Arg Asp Asn Asn Ser His Ser Arg Asp Trp Asp Ile Asn Gly
  65          70          75          80
Gly Ala Cys Asn Thr Leu Thr Asn Asn Glu Gln Asn Gly Pro Lys Leu
          85          90          95
Glu Asn Phe Leu Gly Arg Thr Thr Thr Ile Tyr Asn Thr Asn Glu Thr
  100          105          110
Val Val Asp Gly Asn Gly Asp Cys Gly Gly Gly Asp Gly Gly Gly Gly
  115          120          125
Gly Ser Leu Gly Leu Ser Met Ile Lys Thr Trp Leu Ser Asn His Ser
  130          135          140
Val Ala Asn Ala Asn His Gln Asp Asn Gly Asn Gly Ala Arg Gly Leu
  145          150          155          160
Ser Leu Ser Met Asn Ser Ser Thr Ser Asp Ser Asn Asn Tyr Asn Asn
  165          170          175
Asn Asp Asp Val Val Gln Glu Lys Thr Ile Val Asp Val Val Glu Thr
  180          185          190
Thr Pro Lys Lys Thr Ile Glu Ser Phe Gly Gln Arg Thr Ser Ile Tyr
  195          200          205
Arg Gly Val Thr Arg His Arg Trp Thr Gly Arg Tyr Glu Ala His Leu
  210          215          220
Trp Asp Asn Ser Cys Lys Arg Glu Gly Gln Thr Arg Lys Gly Arg Gln
  225          230          235          240
Val Tyr Leu Gly Gly Tyr Asp Lys Glu Glu Lys Ala Ala Arg Ala Tyr

```

SUBSTITUTE SHEET (RULE 26)

11/13

	245		250		255
Asp Leu Ala Ala Leu Lys Tyr Trp Gly Pro Thr Thr Thr Thr Asn Phe	260		265		270
Pro Leu Ser Glu Tyr Glu Lys Glu Val Glu Glu Met Lys His Met Thr	275		280		285
Arg Gln Glu Tyr Val Ala Ser Leu Arg Arg Lys Ser Ser Gly Phe Ser	290		295		300
Arg Gly Ala Ser Ile Tyr Arg Gly Val Thr Arg His His Gln His Gly	305		310		315
Arg Trp Gln Ala Arg Ile Gly Arg Val Ala Gly Asn Lys Asp Leu Tyr	325		330		335
Leu Gly Thr Phe Gly Thr Gln Glu Glu Ala Ala Glu Ala Tyr Asp Ile	340		345		350
Ala Ala Ile Lys Phe Arg Gly Leu Ser Ala Val Thr Asn Phe Asp Met	355		360		365
Asn Arg Tyr Asn Val Lys Ala Ile Leu Glu Ser Pro Ser Leu Pro Ile	370		375		380
Gly Ser Ser Ala Lys Arg Leu Lys Asp Val Asn Asn Pro Val Pro Ala	385		390		395
Met Met Ile Ser Asn Asn Val Ser Glu Ser Ala Asn Asn Val Ser Gly	405		410		415
Trp Gln Asn Thr Ala Phe Gln His His Gln Gly Met Asp Leu Ser Leu	420		425		430
Leu Gln Gln Gln Gln Glu Arg Tyr Val Gly Tyr Tyr Asn Gly Gly Asn	435		440		445
Leu Ser Thr Glu Ser Thr Arg Val Cys Phe Lys Gln Glu Glu Glu Gln	450		455		460
Gln His Phe Leu Arg Asn Ser Pro Ser His Met Thr Asn Val Asp His	465		470		475
His Ser Ser Thr Ser Asp Asp Ser Val Thr Val Cys Gly Asn Val Val	485		490		495
Ser Tyr Gly Gly Tyr Gln Gly Phe Ala Ile Pro Val Gly Thr Ser Val	500		505		510
Asn Tyr Asp Pro Phe Thr Ala Ala Glu Ile Ala Tyr Asn Ala Arg Asn	515		520		525
His Tyr Tyr Tyr Ala Gln His Gln Gln Gln Gln Ile Gln Gln Ser	530		535		540
Pro Gly Gly Asp Phe Pro Val Ala Ile Ser Asn Asn His Ser Ser Asn	545		550		555
Met Tyr Phe His Gly Glu Gly Gly Gly Glu Gly Ala Pro Thr Phe Ser	565		570		575
Val Trp Asn Asp Thr	580				

12/13

<210> 8
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 8
gaggcagcgg tcggatcgta acagtactct 30

<210> 9
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 9
cataaggaga gagagaaaag cctaaccagt 30

<210> 10
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 10
accaagaact cgtagatc 19

<210> 11
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 11
aacgcatata actaaagatc 20

<210> 12
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 12
ccatggatcc agagacgaag cgaaac 26

<210> 13
<211> 26
<212> DNA
<213> Artificial Sequence

<220>

13 /13

<223> Description of Artificial Sequence: Primer

<400> 13

actccatgga taataactgg ttaggc

26

<210> 14

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 14

aaattctcaa gctttggtcc atcttg

26

2000075330

```

<110> CPRO-DLO
Agriculture and Agri-food Canada

<120> Use of the BNM3 Transcriptional Activator to Control
Plant Embryogenesis and Regeneration Processes

<130> 07-334064

<140>
<141>

<150> EP 99201745.9-2106
<151> 1999-06-02

<160> 14

<170> PatentIn Ver. 2.1

<210> 1
<211> 2014
<212> DNA
<213> Brassica napus

<400> 1
gttcactctct cttctttaag accaaaacct ttttctcttc ctcttcatgc atgaacctta 60
actaagtctct tcttcttttta cctttttacca agaactcgtt agatcactct ctgaactcaa 120
tgaataataaa ctggttaggc tcttctctct ctcttattga acaaaatcac catcgtaagg 180
acgtctactc ttccaccacc acaaccgtcg tagatgtcgc cggagagtag tgtttacgatc 240
ccgacgctgc ctccgatgag tcttcagcca tccaaacatc gtttctctct cctcttgggtg 300
tcgtctgttga tgcgtttcac agagacaaca atagtctact ccgagattgg gacatcaatg 360
gttgtgtcatg caataacatc cacaaacgat agcaagatgg accaaagctt gagaatttcc 420
ttggcgcgac caccacgatt tacaaacaca acgaaaacgt tggagatgga agtgggaagt 480
gctctattagg agggaggagac ggtggtgggt gctcactagg actttcgtat ataaagacat 540
ggcttgagaaa tcaaccocgtg gataatgttg ataatacaga aaatgggcaat gctgcgaaag 600
gcctgtccctc ctcaatgaac tcatctactt cttgtgataa caacaacgac agcaataaca 660
acggttgggtg ccaaggaggag actattgatg atagcgttga agctcacccg aagaaaaacta 720
ttgagaggttt tggacagagg acgtctatat accgcggtgt tacaggatct cggtggacag 780
gaagatatga ggcacattta tgggataata gttgtaaaag agaaggccaa accgcgcaag 840
gaagacaagt ttattttggg gtttatgaca aagaagaaaa agcagctagg gcttatgat 900
tagcgcgact acagtattga ggaaccca ctactactaa cttcccatag agcgaatatg 960
aaaaagaggt agaagagatg aagcatatga caaggccaaga gtatgttgcc tcaactgcga 1020
ggaaaagtag tggtttctct cgtggtgcac ctggttatcg ttgagtaaca agacataccc 1080
aacatggaag atggcaagct aggataggaa gagtgcgccg taacaaagac ctctacttgg 1140
gaactctttgg cacacaagaa gaagctgcag aggcatacga acttcgtgaca caaocgttaa 1200
gaggattaac cgacgttagc aacttcgaca tgaacagata caaocgttaa gcaatctcgc 1260
aaagccctagt tcttctactt ggttagccgc caaaacgatct caaggaggct aactgcgcg 1320
ttccagatgat gattgatgac agtaataacg tttcacgagat tgagaatagt gccagctggt 1380
ggcaaaacgc tgcggttcag catcatcagg gagttagatt gagcttattg caccacaatc 1440
aagagaggtta caatcggttat tattacaagt gaggaaactt gtcttcctagg agtgcctagg 1500
cttgtttcaa acaagaggat gatcaacacc atttcttgag caaacgcgag agcctcatga 1560
ctaattatcga tcatcaaaat tctgtttcgg atgatctcgt tactgtgttg ggaaatgcta 1620
ttggttatgg tggttatcaa ggaatttcag ccccggttaa ctgcgattgc tacgtccta 1680

```

```
<210> 2
<211> 579
<212> PRT
<213> Brassica napus
```

```

400> 2
Met Asn Asn Asn Trp Leu Gly Phe Ser Leu Ser Pro Tyr Glu Gln Asn
  1          5          10          15

His His Arg Lys Asp Val Tyr Ser Ser Thr Thr Thr Thr Val Val Asp
      20          25          30

Val Ala Gly Glu Tyr Cys Tyr Asp Pro Thr Ala Ala Ser Asp Glu Ser
      35          40          45

Ser Ala Ile Gln Thr Ser Phe Pro Ser Pro Phe Gly Val Val Val Asp
      50          55          60

Ala Phe Thr Arg Asp Asn Asn Ser His Ser Arg Asp Trp Asp Ile Asn
      65          70          75          80

Gly Cys Ala Cys Asn Asn Ile His Asn Asp Glu Gln Asp Gly Pro Lys
      85          90          95

Leu Glu Asn Phe Leu Gly Arg Thr Thr Thr Ile Tyr Asn Thr Asn Glu
      100          105          110

Asn Val Gly Asp Gly Ser Gly Ser Gly Cys Tyr Gly Gly Gly Asp Gly
      115          120          125

Gly Gly Gly Ser Leu Gly Leu Ser Met Ile Lys Thr Trp Leu Arg Asn
      130          135          140

Gln Pro Val Asp Asn Val Asp Asn Gln Glu Asn Gly Asn Ala Ala Lys
      145          150          155          160

Gly Leu Ser Leu Ser Met Asn Ser Ser Thr Ser Cys Asp Asn Asn Asn
      165          170          175

Asp Ser Asn Asn Asn Val Val Ala Gln Gly Lys Thr Ile Asp Asp Ser
      180          185          190

Val Glu Ala Thr Pro Lys Lys Thr Ile Glu Ser Phe Gly Gln Arg Thr
      195          200          205

Ser Ile Tyr Arg Gly Val Thr Arg His Arg Trp Thr Gly Arg Tyr Glu
      210          215          220

```

Ala	His	Leu	Trp	Asp	Asn	Asn	Ser	Cys	Lys	Arg	Glu	Gly	Gln	Thr	Arg	Lys
225					230						235					240
Gly	Arg	Gln	Val	Tyr	Leu	Gly	Gly	Tyr	Asp	Lys	Glu	Glu	Lys	Ala	Ala	
				245					250					255		
Arg	Ala	Tyr	Asp	Leu	Ala	Ala	Leu	Lys	Tyr	Trp	Gly	Thr	Thr	Thr	Thr	
			260					265					270			
Thr	Asn	Phe	Pro	Met	Ser	Glu	Tyr	Glu	Lys	Glu	Val	Glu	Glu	Met	Lys	
		275					280					285				
His	Met	Thr	Arg	Gln	Glu	Tyr	Val	Ala	Ser	Leu	Arg	Arg	Lys	Ser	Ser	
	290					295					300					
Gly	Phe	Ser	Arg	Gly	Ala	Ser	Ile	Tyr	Arg	Gly	Val	Thr	Arg	His	His	
305					310					315				320		
Gln	His	Gly	Arg	Trp	Gln	Ala	Arg	Ile	Gly	Arg	Val	Ala	Gly	Asn	Lys	
				325					330					335		
Asp	Leu	Tyr	Leu	Gly	Thr	Phe	Gly	Thr	Gln	Glu	Glu	Ala	Ala	Glu	Ala	
			340					345					350			
Tyr	Asp	Ile	Ala	Ala	Ile	Lys	Phe	Arg	Gly	Leu	Thr	Ala	Val	Thr	Asn	
		355					360					365				
Phe	Asp	Met	Asn	Arg	Tyr	Asn	Val	Lys	Ala	Ile	Leu	Glu	Ser	Pro	Ser	
	370					375					380					
Leu	Pro	Ile	Gly	Ser	Ala	Ala	Lys	Arg	Leu	Lys	Glu	Ala	Asn	Arg	Pro	
385					390					395				400		
Val	Pro	Ser	Met	Met	Met	Ile	Ser	Asn	Asn	Val	Ser	Glu	Ser	Glu	Asn	
				405					410					415		
Ser	Ala	Ser	Gly	Trp	Gln	Asn	Ala	Ala	Val	Gln	His	His	Gln	Gly	Val	
			420					425					430			
Asp	Leu	Ser	Leu	Leu	His	Gln	His	Gln	Glu	Arg	Tyr	Asn	Gly	Tyr	Tyr	
		435				440						445				
Tyr	Asn	Gly	Gly	Asn	Leu	Ser	Ser	Glu	Ser	Ala	Arg	Ala	Cys	Phe	Lys	
	450					455					460					
Gln	Glu	Asp	Asp	Gln	His	His	Phe	Leu	Ser	Asn	Thr	Gln	Ser	Leu	Met	
465				470						475				480		
Thr	Asn	Ile	Asp	His	Gln	Ser	Ser	Val	Ser	Asp	Asp	Ser	Val	Thr	Val	
				485					490					495		
Cys	Gly	Asn	Val	Val	Gly	Tyr	Gly	Gly	Tyr	Gln	Gly	Phe	Ala	Ala	Pro	
			500					505					510			
Val	Asn	Cys	Asp	Ala	Tyr	Ala	Ala	Ser	Glu	Phe	Asp	Tyr	Asn	Ala	Arg	

525

Asn Asp Asn

[illegible]

<210> 4
 <211> 579
 <212> PRT
 <213> Brassica napus

<400> 4
 Met Asn Asn Asn Trp Leu Gly Phe Ser Leu Ser Pro Tyr Glu Gln Asn
 1 5 10 15
 His His Arg Lys Asp Val Tyr Ser Ser Thr Thr Thr Thr Val Val Asp
 20 25 30
 Val Ala Gly Glu Tyr Cys Tyr Asp Pro Thr Ala Ala Ser Asp Glu Ser
 35 40 45
 Ser Ala Ile Gln Thr Ser Phe Pro Ser Pro Phe Gly Val Val Val Asp
 50 55 60
 Ala Phe Thr Arg Asp Asn Asn Ser His Ser Arg Asp Trp Asp Ile Asn
 65 70 75 80
 Gly Cys Ala Cys Asn Asn Ile His Asn Asp Glu Gln Asp Gly Pro Lys
 85 90 95
 Leu Glu Asn Phe Leu Gly Arg Thr Thr Thr Ile Tyr Asn Thr Asn Glu
 100 105 110
 Asn Val Gly Asp Gly Ser Gly Ser Gly Cys Tyr Gly Gly Gly Asp Gly
 115 120 125
 Gly Gly Gly Ser Leu Gly Leu Ser Met Ile Lys Thr Trp Leu Arg Asn
 130 135 140
 Gln Pro Val Asp Asn Val Asp Asn Gln Glu Asn Gly Asn Ala Ala Lys
 145 150 155 160
 Gly Leu Ser Leu Ser Met Asn Ser Ser Thr Ser Cys Asp Asn Asn Asn
 165 170 175
 Asp Ser Asn Asn Asn Val Val Ala Gln Gly Lys Thr Ile Asp Asp Ser
 180 185 190
 Val Glu Ala Thr Pro Lys Lys Thr Ile Glu Ser Phe Gly Gln Arg Thr
 195 200 205
 Ser Ile Tyr Arg Gly Val Thr Arg His Arg Trp Thr Gly Arg Tyr Glu
 210 215 220
 Ala His Leu Trp Asp Asn Ser Cys Lys Arg Glu Gly Gln Thr Arg Lys
 225 230 235 240
 Gly Arg Gln Val Tyr Leu Gly Gly Tyr Asp Lys Glu Glu Lys Ala Ala
 245 250 255

2000-1-19-1999

Arg	Ala	Tyr	Asp	Leu	Ala	Ala	Leu	Lys	Tyr	Trp	Gly	Thr	Thr	Thr	Thr		
			260						265						270		
Thr	Asn	Phe	Pro	Met	Ser	Glu	Tyr	Glu	Lys	Glu	Val	Glu	Glu	Met	Lys		
		275					280						285				
His	Met	Thr	Arg	Gln	Glu	Tyr	Val	Ala	Ser	Leu	Arg	Arg	Lys	Ser	Ser		
		290				295					300						
Gly	Phe	Ser	Arg	Gly	Ala	Ser	Ile	Tyr	Arg	Gly	Val	Thr	Arg	His	His		
305					310						315				320		
Gln	His	Gly	Arg	Trp	Gln	Ala	Arg	Ile	Gly	Arg	Val	Ala	Gly	Asn	Lys		
				325						330					335		
Asp	Leu	Tyr	Leu	Gly	Thr	Phe	Gly	Thr	Gln	Glu	Glu	Ala	Ala	Glu	Ala		
			340					345					350				
Tyr	Asp	Ile	Ala	Ala	Ile	Lys	Phe	Arg	Gly	Leu	Thr	Ala	Val	Thr	Asn		
		355					360						365				
Phe	Asp	Met	Asn	Arg	Tyr	Asn	Val	Lys	Ala	Ile	Leu	Glu	Ser	Pro	Ser		
		370				375						380					
Leu	Pro	Ile	Gly	Ser	Ala	Ala	Lys	Arg	Leu	Lys	Glu	Ala	Asn	Arg	Pro		
385					390					395					400		
Val	Pro	Ser	Met	Met	Met	Ile	Ser	Asn	Asn	Val	Ser	Glu	Ser	Glu	Asn		
				405					410						415		
Ser	Ala	Ser	Gly	Trp	Gln	Asn	Ala	Ala	Val	Gln	His	His	Gln	Gly	Val		
			420					425					430				
Asp	Leu	Ser	Leu	Leu	His	Gln	His	Gln	Glu	Arg	Tyr	Asn	Gly	Tyr	Tyr		
		435				440						445					
Tyr	Asn	Gly	Gly	Asn	Leu	Ser	Ser	Glu	Ser	Ala	Arg	Ala	Cys	Phe	Lys		
	450				455						460						
Gln	Glu	Asp	Asp	Gln	His	His	Phe	Leu	Ser	Asn	Thr	Gln	Ser	Leu	Met		
465				470						475					480		
Thr	Asn	Ile	Asp	His	Gln	Ser	Ser	Val	Ser	Asp	Asp	Ser	Val	Thr	Val		
				485					490						495		
Cys	Gly	Asn	Val	Val	Gly	Tyr	Gly	Gly	Tyr	Gln	Gly	Phe	Ala	Ala	Pro		
		500						505					510				
Val	Asn	Cys	Asp	Ala	Tyr	Ala	Ala	Ser	Glu	Phe	Asp	Tyr	Asn	Ala	Arg		
		515					520					525					
Asn	His	Tyr	Tyr	Phe	Ala	Gln	Gln	Gln	Gln	Thr	Gln	Gln	Ser	Pro	Gly		
	530					535						540					
Gly	Asp	Phe	Pro	Ala	Ala	Met	Thr	Asn	Asn	Val	Gly	Ser	Asn	Met	Tyr		

7/15

545

550

555

560

Tyr His Gly Glu Gly Gly Glu Val Ala Pro Thr Phe Thr Val Trp
565 570 575

Asn Asp Asn

<210> 5

<211> 4873

<212> DNA

<213> Brassica napus

<220>

<221> intron

<222> (1846)..(2298)

<220>

<221> intron

<222> (2720)..(2952)

<220>

<221> intron

<222> (3036)..(3160)

<220>

<221> intron

<222> (3170)..(3314)

<220>

<221> intron

<222> (3404)..(3553)

<220>

<221> intron

<222> (3628)..(3797)

<220>

<221> intron

<222> (3849)..(3961)

<220>

<221> intron

<222> (4039)..(4148)

<220>

<221> misc_feature

<222> (1620)..(1622)

<223> start codon

<220>

<221> misc_feature

<222> (4856)..(4858)

<223> stop codon

20250401 14:50:00

[illegible]

tttttgtgta tacattcatt atataatcaa aatagcgata cगततततcat tcaattgttg 3300
 tctatatcat gcaggagggt atgacaaaga agaaaaagca gctaggggctt atgatttagc 3360
 cgcactcaa ttttggggaa ccaccactac tactaacttc cccgtaagtc aatcaatgtt 3420
 gtacaagatt tcataactta gaaccaattt tattcttttt ttataaagatg ctattatctt 3480
 attattaatt gccattgttta tatcggttaca tttattacaa taaaaagtac ttttgggttg 3540
 atataatatt tagatgagcg aatatgaaaa agaggtagaa gagatgaagc acatgacaag 3600
 gcaagagtat gttgcctcac tgcgcaggta tataatggaa cttctgatat tattgcatat 3660
 ggcatctatt attatacatg tatattagta ttttataat agaacccatc acgctcacgt 3720
 ttatatttaa aaatatgtcc gtattcacgt cagattatca gcatacacct atataataa 3780
 gacattaaaa tatgcaggaa aagtagtggt ttctctcgtg gtgcatacat ttatcgtgga 3840
 gtaacaaggt attcatacag agagaacgaa tctatatttg ttacgtacat atatatataa 3900
 aaatataatt ataagatatt acattttata ttatgaatat ttcttctaatt ggggtccaaaa 3960
 gacatcacca acatggaaga tggcaagcta ggataggaag agtcgcgggt aacaaagacc 4020
 tctacttggg aacttttgggt acgttttagtc ttctcttact aaacttcaca atcaaatcta 4080
 taacaaaaaga tatcaactaa aaactacaac atatatctaa gtaagctgta catatattat 4140
 atatgaagtc acacaagaag aagctgcaga ggcatcacgac attgcggcca tcaaatcag 4200
 aggattaaac gcagtgacta acttcgacat gaacagatac aacgttaaag caatcctcga 4260
 aagccctagt cttctcattg gtacgcgcgc aaaacgtctc aaggaggcta accgtccggt 4320
 tccaagtatt atgatgatca gtaataacgt ttccagagagt gagaatagtg ctacgggttg 4380
 gcaaaacgct gcgggtcagc atcatcaggg agtagatttg agcttattgc accaacaatca 4440
 agagaggtac aatggttatt attacaatgg agggaaacttg tcttcggaga gtgctagggc 4500
 ttgtttcaaa caagaggatg atcaacacca tttcttgagc aacacgcaga gcctcatgac 4560
 taatatcgat catcaaagtt ctgtttcgga tgattcgggt actgtttgtg gaaatgttgt 4620
 tgggtatggt ggttatcaag gatttcgagc cccggttaac tgcgatgcct acgctgctag 4680
 tgaagtttga tataacgcaa gaaaccatta ttactttgct cagcagcagc agaccagca 4740
 gtccgcaggt ggagattttc ccgcggcaat gacgaataat gttggctcta atatgtatta 4800
 ccattgggaa ggtgtgggag aagttgctcc aacatttaca gtttggaaac acaattagaa 4860
 aaaatagtta aag 4873

<210> 6

<211> 5151

<212> DNA

<213> Arabidopsis thaliana

<220>

<221> intron

<222> (2249)..(2578)

<220>

<221> intron

<222> (2994)..(3220)

<220>

<221> intron

<222> (3304)..(3420)

<220>

<221> intron

<222> (3429)..(3521)

<220>

<221> intron

<222> (3611)..(3770)

[illegible]

5151

```

400> 7
Met Asn Asn Trp Leu Gly Phe Ser Leu Ser Pro His Asp Gln Asn His
  1                      5                      10                      15

His Arg Thr Asp Val Asp Ser Ser Thr Thr Arg Thr Ala Val Asp Val
                      20                      25                      30

Ala Gly Gly Tyr Cys Phe Asp Leu Ala Ala Pro Ser Asp Glu Ser Ser
  35                      40                      45

Ala Val Gln Thr Ser Phe Leu Ser Pro Phe Gly Val Thr Leu Glu Ala
  50                      55                      60

Phe Thr Arg Asp Asn Asn Ser His Ser Arg Asp Trp Asp Ile Asn Gly
  65                      70                      75                      80

Gly Ala Cys Asn Thr Leu Thr Asn Asn Glu Gln Asn Gly Pro Lys Leu
                      85                      90                      95

Glu Asn Phe Leu Gly Arg Thr Thr Thr Ile Tyr Asn Thr Asn Glu Thr
  100                      105                      110

Val Val Asp Gly Asn Gly Asp Cys Gly Gly Gly Asp Gly Gly Gly Gly
  115                      120                      125

Gly Ser Leu Gly Leu Ser Met Ile Lys Thr Trp Leu Ser Asn His Ser
  130                      135                      140

Val Ala Asn Ala Asn His Gln Asp Asn Gly Asn Gly Ala Arg Gly Leu
  145                      150                      155                      160

Ser Leu Ser Met Asn Ser Ser Thr Ser Asp Ser Asn Asn Tyr Asn Asn
                      165                      170                      175

Asn Asp Asp Val Val Gln Glu Lys Thr Ile Val Asp Val Val Glu Thr
  180                      185                      190

Thr Pro Lys Lys Thr Ile Glu Ser Phe Gly Gln Arg Thr Ser Ile Tyr
  195                      200                      205

Arg Gly Val Thr Arg His Arg Trp Thr Gly Arg Tyr Glu Ala His Leu
  210                      215                      220

Trp Asp Asn Ser Cys Lys Arg Glu Gly Gln Thr Arg Lys Gly Arg Gln
  225                      230                      235                      240

Val Tyr Leu Gly Gly Tyr Asp Lys Glu Glu Lys Ala Ala Arg Ala Tyr
  245                      250                      255

Asp Leu Ala Ala Leu Lys Tyr Trp Gly Pro Thr Thr Thr Thr Asn Phe

```

270

Pro Gly Gly Asp Phe Pro Val Ala Ile Ser Asn Asn His Ser Ser Asn
545 550 555 560

Met Tyr Phe His Gly Glu Gly Gly Gly Glu Gly Ala Pro Thr Phe Ser
 565 570 575

Val Trp Asn Asp Thr
 580

<210> 8
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 8
 gaggcagcgg tcggatcgta acagtactct 30

<210> 9
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 9
 cataaggaga gagagaaaag cctaaccagt 30

<210> 10
 <211> 19
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 10
 accaagaact cggttagatc 19

<210> 11
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 11
 aacgcatata actaaagatc 20

<210> 12
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 12
ccatggatcc agagacgaag cgaaac

26

<210> 13
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 13
actccatgga taataactgg ttaggc

26

<210> 14
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 14
aaattctcaa gctttgtcc atcttg

26